

表 875 1セッションあたりの置換液量(L) 透析歴(年)別 (オフラインHDF 透析回数週3回 後希釈)

透析歴	1未満	1～	5～	10～	15～	20～	合計	記載なし	総計	平均	標準偏差
2年未満		37	171	48	1	1	258	9	267	7.20	2.65
(%)		(14.3)	(66.3)	(18.6)	(0.4)	(0.4)	(100.0)				
2年～		58	309	156	1	1	525	24	549	7.64	2.32
(%)		(11.0)	(58.9)	(29.7)	(0.2)	(0.2)	(100.0)				
5年～		69	411	264			744	29	773	7.93	2.22
(%)		(9.3)	(55.2)	(35.5)			(100.0)				
10年～		38	271	211	2		522	27	549	8.16	2.09
(%)		(7.3)	(51.9)	(40.4)	(0.4)		(100.0)				
15年～		26	197	119	1	1	344	10	354	8.09	2.11
(%)		(7.6)	(57.3)	(34.6)	(0.3)	(0.3)	(100.0)				
20年～		14	112	138	1		265	13	278	8.52	2.08
(%)		(5.3)	(42.3)	(52.1)	(0.4)		(100.0)				
25年～		6	85	101			192	12	204	8.60	1.86
(%)		(3.1)	(44.3)	(52.6)			(100.0)				
30年～		6	59	71			136	6	142	8.64	2.00
(%)		(4.4)	(43.4)	(52.2)			(100.0)				
35年～		5	43	39	1		88	5	93	8.40	2.45
(%)		(5.7)	(48.9)	(44.3)	(1.1)		(100.0)				
40年～		4	17	16			37	2	39	7.89	2.53
(%)		(10.8)	(45.9)	(43.2)			(100.0)				
合計		263	1,675	1,163	7	3	3,111	137	3,248	8.01	2.24
(%)		(8.5)	(53.8)	(37.4)	(0.2)	(0.1)	(100.0)				
不明			1				1	1	2	8.00	
(%)			(100.0)				(100.0)				
記載なし											
(%)											
総計		263	1,676	1,163	7	3	3,112	138	3,250	8.01	2.24
(%)		(8.5)	(53.9)	(37.4)	(0.2)	(0.1)	(100.0)				
平均		9.69	11.51	14.66	14.71	7.33	12.54	13.15	12.56		
標準偏差		9.30	9.70	10.31	12.16	10.21	10.05	10.60	10.07		

数値下のかっこ内は行方向の合計に対する%です。

##D1932##

(C)Japanese Society for Dialysis Therapy

2016年調査