

表 876 1セッションあたりの置換液量(L) 原疾患別 (オフラインHDF 透析回数週3回 後希釈)

原疾患	1未満	1~	5~	10~	15~	20~	合計	記載なし	総計	平均	標準偏差
慢性系球体腎炎 (%)		92 (7.6)	592 (48.8)	525 (43.3)	4 (0.3)		1,213 (100.0)	70	1,283	8.21	2.15
慢性腎孟腎炎 (%)			24 (70.6)	10 (29.4)			34 (100.0)	1	35	7.88	1.63
急速進行性系球体腎炎 (%)	2 (8.7)	14 (60.9)	7 (30.4)				23 (100.0)	1	24	7.65	2.14
妊娠腎/妊娠中毒症 (%)	2 (5.6)	19 (52.8)	15 (41.7)				36 (100.0)	1	37	8.11	2.29
その他の分類不能の腎炎 (%)	1 (5.9)	6 (35.3)	10 (58.8)				17 (100.0)	1	18	8.65	2.26
多発性囊胞腎 (%)	11 (9.5)	62 (53.4)	42 (36.2)		1 (0.9)		116 (100.0)	2	118	8.18	2.97
腎硬化症 (%)	22 (11.5)	124 (64.9)	45 (23.6)				191 (100.0)	7	198	7.47	2.15
悪性高血圧 (%)	1 (5.6)	11 (61.1)	6 (33.3)				18 (100.0)	2	20	8.17	1.76
糖尿病性腎症 (%)	87 (8.7)	564 (56.3)	348 (34.8)	1 (0.1)	1 (0.1)		1,001 (100.0)	40	1,041	7.87	2.24
SLE腎炎 (%)	4 (10.5)	19 (50.0)	15 (39.5)				38 (100.0)	2	40	7.84	2.30
アミロイド腎 (%)		4 (57.1)	3 (42.9)				7 (100.0)	1	8	8.57	1.51
痛風腎 (%)		4 (33.3)	8 (66.7)				12 (100.0)		12	9.33	1.50
先天性代謝異常による腎不全 (%)		1 (50.0)	1 (50.0)				2 (100.0)		2	9.00	1.41
腎・尿路結核 (%)											
腎・尿路結石 (%)	1 (14.3)	5 (71.4)	1 (14.3)				7 (100.0)		7	7.57	1.99
腎・尿路腫瘍 (%)		1 (9.1)	5 (45.5)	5 (45.5)			11 (100.0)		11	8.27	2.53
閉塞性尿路障害 (%)	1 (14.3)	4 (57.1)	2 (28.6)				7 (100.0)		7	7.43	2.23
骨髄腫 (%)		4 (80.0)	1 (20.0)				5 (100.0)		5	6.80	1.79
腎形成不全 (%)		6 (54.5)	4 (36.4)	1 (9.1)			11 (100.0)		11	9.73	2.61
不明 (%)	23 (9.1)	154 (60.9)	75 (29.6)		1 (0.4)		253 (100.0)	6	259	7.86	2.18
再導入 (%)	4 (9.3)	22 (51.2)	16 (37.2)	1 (2.3)			43 (100.0)	1	44	8.12	2.52
その他 (%)	11 (16.4)	32 (47.8)	24 (35.8)				67 (100.0)	3	70	7.82	2.69
合計 (%)		263 (8.5)	1,676 (53.9)	1,163 (37.4)	7 (0.2)	3 (0.1)	3,112 (100.0)	138	3,250	8.01	2.24
記載なし (%)											
総計 (%)		263 (8.5)	1,676 (53.9)	1,163 (37.4)	7 (0.2)	3 (0.1)	3,112 (100.0)	138	3,250	8.01	2.24

数値下のカッコ内は行方向の合計に対する%です。

##D1933##

(C)Japanese Society for Dialysis Therapy

2016年調査