

表 894 身長(cm) 原疾患別 (透析患者全体)

原疾患	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	55	191	795	2,959	7,353	12,286	13,917	15,745	14,349	9,432	3,398	842	125	14	81,461	10,322	91,783	159.98	9.44
(%)	(0.1)	(0.2)	(1.0)	(3.6)	(9.0)	(15.1)	(17.1)	(19.3)	(17.6)	(11.6)	(4.2)	(1.0)	(0.2)	(0.0)	(100.0)				
慢性腎盂腎炎	2	13	34	147	334	472	474	445	341	210	89	26		1	2,588	308	2,896	157.77	9.68
(%)	(0.1)	(0.5)	(1.3)	(5.7)	(12.9)	(18.2)	(18.3)	(17.2)	(13.2)	(8.1)	(3.4)	(1.0)		(0.0)	(100.0)				
急速進行性糸球体腎炎	2	4	38	130	244	368	389	408	301	232	71	22	5		2,214	324	2,538	158.40	9.79
(%)	(0.1)	(0.2)	(1.7)	(5.9)	(11.0)	(16.6)	(17.6)	(18.4)	(13.6)	(10.5)	(3.2)	(1.0)	(0.2)		(100.0)				
妊娠腎/妊娠中毒症	2	2	16	98	278	430	363	148	38	5	1				1,381	102	1,483	152.81	6.23
(%)	(0.1)	(0.1)	(1.2)	(7.1)	(20.1)	(31.1)	(26.3)	(10.7)	(2.8)	(0.4)	(0.1)				(100.0)				
その他の分類不能の腎炎	5	1	19	54	116	185	218	221	244	166	64	16	2		1,311	143	1,454	160.07	10.28
(%)	(0.4)	(0.1)	(1.4)	(4.1)	(8.8)	(14.1)	(16.6)	(16.9)	(18.6)	(12.7)	(4.9)	(1.2)	(0.2)		(100.0)				
多発性嚢胞腎	9	8	51	210	722	1,452	1,804	1,772	1,688	1,488	763	226	50	6	10,249	1,222	11,471	161.89	9.84
(%)	(0.1)	(0.1)	(0.5)	(2.0)	(7.0)	(14.2)	(17.6)	(17.3)	(16.5)	(14.5)	(7.4)	(2.2)	(0.5)	(0.1)	(100.0)				
腎硬化症	24	135	506	1,567	2,875	3,957	4,733	5,667	4,569	2,529	769	188	25	3	27,547	4,103	31,650	158.54	9.61
(%)	(0.1)	(0.5)	(1.8)	(5.7)	(10.4)	(14.4)	(17.2)	(20.6)	(16.6)	(9.2)	(2.8)	(0.7)	(0.1)	(0.0)	(100.0)				
悪性高血圧		7	28	71	183	275	347	479	471	343	155	43	6	2	2,410	311	2,721	161.70	9.73
(%)		(0.3)	(1.2)	(2.9)	(7.6)	(11.4)	(14.4)	(19.9)	(19.5)	(14.2)	(6.4)	(1.8)	(0.2)	(0.1)	(100.0)				
糖尿病性腎症	100	159	734	3,146	7,622	12,933	16,192	22,598	22,069	14,932	5,757	1,652	263	50	108,207	15,450	123,657	161.58	9.38
(%)	(0.1)	(0.1)	(0.7)	(2.9)	(7.0)	(12.0)	(15.0)	(20.9)	(20.4)	(13.8)	(5.3)	(1.5)	(0.2)	(0.0)	(100.0)				
SLE腎炎	1	5	27	91	237	436	430	328	216	111	44	16	2		1,944	278	2,222	156.91	8.76
(%)	(0.1)	(0.3)	(1.4)	(4.7)	(12.2)	(22.4)	(22.1)	(16.9)	(11.1)	(5.7)	(2.3)	(0.8)	(0.1)		(100.0)				
アミロイド腎	1	1	8	30	44	78	72	55	43	37	3	1	1		374	64	438	156.57	9.70
(%)	(0.3)	(0.3)	(2.1)	(8.0)	(11.8)	(20.9)	(19.3)	(14.7)	(11.5)	(9.9)	(0.8)	(0.3)	(0.3)		(100.0)				
痛風腎			2	16	32	78	140	232	243	143	43	9	4		942	114	1,056	163.16	7.85
(%)			(0.2)	(1.7)	(3.4)	(8.3)	(14.9)	(24.6)	(25.8)	(15.2)	(4.6)	(1.0)	(0.4)		(100.0)				
先天性代謝異常による腎不全	3	3	11	4	29	33	52	43	40	24	12	2	1		257	24	281	158.26	11.81
(%)	(1.2)	(1.2)	(4.3)	(1.6)	(11.3)	(12.8)	(20.2)	(16.7)	(15.6)	(9.3)	(4.7)	(0.8)	(0.4)		(100.0)				
腎・尿路結核	1		4	5	26	28	28	29	17	7	2				147	21	168	156.23	8.74
(%)	(0.7)		(2.7)	(3.4)	(17.7)	(19.0)	(19.0)	(19.7)	(11.6)	(4.8)	(1.4)				(100.0)				
腎・尿路結石	3	3	6	20	44	73	108	113	78	49	18	2			517	61	578	159.02	9.23
(%)	(0.6)	(0.6)	(1.2)	(3.9)	(8.5)	(14.1)	(20.9)	(21.9)	(15.1)	(9.5)	(3.5)	(0.4)			(100.0)				
腎・尿路腫瘍	1	2	8	20	57	80	122	202	205	101	39	7	1		845	114	959	161.62	8.90
(%)	(0.1)	(0.2)	(0.9)	(2.4)	(6.7)	(9.5)	(14.4)	(23.9)	(24.3)	(12.0)	(4.6)	(0.8)	(0.1)		(100.0)				
閉塞性尿路障害	4	5	5	32	70	96	113	117	116	71	24	5	2	1	661	82	743	159.02	10.60
(%)	(0.6)	(0.8)	(0.8)	(4.8)	(10.6)	(14.5)	(17.1)	(17.7)	(17.5)	(10.7)	(3.6)	(0.8)	(0.3)	(0.2)	(100.0)				
骨髄腫	1	1	2	16	25	45	55	55	40	28	8	3			279	44	323	158.74	9.44
(%)	(0.4)	(0.4)	(0.7)	(5.7)	(9.0)	(16.1)	(19.7)	(19.7)	(14.3)	(10.0)	(2.9)	(1.1)			(100.0)				
腎形成不全	28	14	22	39	56	61	94	113	83	65	18	9			602	62	664	155.48	16.83
(%)	(4.7)	(2.3)	(3.7)	(6.5)	(9.3)	(10.1)	(15.6)	(18.8)	(13.8)	(10.8)	(3.0)	(1.5)			(100.0)				
不明	37	96	393	1,307	2,747	4,014	4,422	5,222	4,448	2,638	961	247	43	3	26,578	4,834	31,412	159.02	9.73
(%)	(0.1)	(0.4)	(1.5)	(4.9)	(10.3)	(15.1)	(16.6)	(19.6)	(16.7)	(9.9)	(3.6)	(0.9)	(0.2)	(0.0)	(100.0)				
再導入	3	6	12	38	145	221	299	396	390	263	116	31	4	1	1,925	249	2,174	161.80	9.37
(%)	(0.2)	(0.3)	(0.6)	(2.0)	(7.5)	(11.5)	(15.5)	(20.6)	(20.3)	(13.7)	(6.0)	(1.6)	(0.2)	(0.1)	(100.0)				
その他	28	28	91	345	682	998	1,119	1,334	1,259	793	302	88	18	1	7,086	1,352	8,438	159.56	10.44
(%)	(0.4)	(0.4)	(1.3)	(4.9)	(9.6)	(14.1)	(15.8)	(18.8)	(17.8)	(11.2)	(4.3)	(1.2)	(0.3)	(0.0)	(100.0)				
合計	310	684	2,812	10,345	23,921	38,599	45,491	55,722	51,248	33,667	12,657	3,435	552	82	279,525	39,584	319,109	160.36	9.62
(%)	(0.1)	(0.2)	(1.0)	(3.7)	(8.6)	(13.8)	(16.3)	(19.9)	(18.3)	(12.0)	(4.5)	(1.2)	(0.2)	(0.0)	(100.0)				
記載なし																			
(%)																			
総計	310	684	2,812	10,345	23,921	38,599	45,491	55,722	51,248	33,667	12,657	3,435	552	82	279,525	39,584	319,109	160.36	9.62
(%)	(0.1)	(0.2)	(1.0)	(3.7)	(8.6)	(13.8)	(16.3)	(19.9)	(18.3)	(12.0)	(4.5)	(1.2)	(0.2)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D2016##

(C)Japanese Society for Dialysis Therapy

2016年調査