

表 914 透析後体重(kg) 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	20未満	20~	30~	40~	50~	60~	70~	80~	90~	100~	110~	120~	合計	記載なし	総計	平均	標準偏差
慢性系球体腎炎		253	5,862	16,941	18,049	9,802	3,206	992	307	119	51	27	55,609	1,563	57,172	53.50	12.02
(%)		(0.5)	(10.5)	(30.5)	(32.5)	(17.6)	(5.8)	(1.8)	(0.6)	(0.2)	(0.1)	(0.0)	(100.0)				
慢性腎孟腎炎		13	288	604	518	266	93	30	14	5	1	1	1,833	35	1,868	51.86	12.56
(%)		(0.7)	(15.7)	(33.0)	(28.3)	(14.5)	(5.1)	(1.6)	(0.8)	(0.3)	(0.1)	(0.1)	(100.0)				
急速進行性系球体腎炎		12	285	538	480	222	59	21	4				1,621	66	1,687	50.48	11.08
(%)		(0.7)	(17.6)	(33.2)	(29.6)	(13.7)	(3.6)	(1.3)	(0.2)				(100.0)				
妊娠腎/妊娠中毒症		4	145	346	232	76	19	6	3	1			832	17	849	48.69	10.12
(%)		(0.5)	(17.4)	(41.6)	(27.9)	(9.1)	(2.3)	(0.7)	(0.4)	(0.1)			(100.0)				
その他の分類不能の腎炎		8	84	266	274	153	50	29	5	3	3	2	877	25	902	54.30	13.77
(%)		(0.9)	(9.6)	(30.3)	(31.2)	(17.4)	(5.7)	(3.3)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
多発性囊胞腎		11	452	1,972	2,428	1,572	651	186	56	18	7	5	7,358	190	7,548	56.18	12.14
(%)		(0.1)	(6.1)	(26.8)	(33.0)	(21.4)	(8.8)	(2.5)	(0.8)	(0.2)	(0.1)	(0.1)	(100.0)				
腎硬化症		117	2,304	6,338	6,616	3,684	1,239	426	139	67	22	14	20,966	552	21,518	53.61	12.48
(%)		(0.6)	(11.0)	(30.2)	(31.6)	(17.6)	(5.9)	(2.0)	(0.7)	(0.3)	(0.1)	(0.1)	(100.0)				
悪性高血圧		6	142	395	491	350	192	94	31	12	2	6	1,721	56	1,777	58.09	14.98
(%)		(0.3)	(8.3)	(23.0)	(28.5)	(20.3)	(11.2)	(5.5)	(1.8)	(0.7)	(0.1)	(0.3)	(100.0)				
糖尿病性腎症		128	4,169	17,295	26,132	19,136	9,106	3,793	1,428	537	187	108	82,019	2,449	84,468	58.97	13.71
(%)		(0.2)	(5.1)	(21.1)	(31.9)	(23.3)	(11.1)	(4.6)	(1.7)	(0.7)	(0.2)	(0.1)	(100.0)				
SLE腎炎		6	237	542	338	138	43	16	6				1,326	44	1,370	49.34	10.94
(%)		(0.5)	(17.9)	(40.9)	(25.5)	(10.4)	(3.2)	(1.2)	(0.5)				(100.0)				
アミロイド腎		8	59	105	62	29	11	1	1				276	4	280	48.34	11.36
(%)		(2.9)	(21.4)	(38.0)	(22.5)	(10.5)	(4.0)	(0.4)	(0.4)				(100.0)				
痛風腎		3	35	137	261	172	51	13	3	4			679	21	700	56.76	11.14
(%)		(0.4)	(5.2)	(20.2)	(38.4)	(25.3)	(7.5)	(1.9)	(0.4)	(0.6)			(100.0)				
先天性代謝異常による腎不全		5	24	54	42	30	7	1	1				164	5	169	51.12	12.11
(%)		(3.0)	(14.6)	(32.9)	(25.6)	(18.3)	(4.3)	(0.6)	(0.6)				(100.0)				
腎・尿路結核		1	23	31	34	10	2						101	5	106	48.61	9.93
(%)		(1.0)	(22.8)	(30.7)	(33.7)	(9.9)	(2.0)						(100.0)				
腎・尿路結石		4	36	107	125	68	24	4	3		1		372	12	384	53.72	12.13
(%)		(1.1)	(9.7)	(28.8)	(33.6)	(18.3)	(6.5)	(1.1)	(0.8)		(0.3)		(100.0)				
腎・尿路腫瘍		41	160	233	128	50	16	5					633	11	644	55.64	11.47
(%)		(6.5)	(25.3)	(36.8)	(20.2)	(7.9)	(2.5)	(0.8)					(100.0)				
閉塞性尿路障害		2	57	122	156	80	24	7	2	1			451	14	465	53.30	11.69
(%)		(0.4)	(12.6)	(27.1)	(34.6)	(17.7)	(5.3)	(1.6)	(0.4)	(0.2)			(100.0)				
骨髄腫		1	28	69	68	28	9	2					205	8	213	51.54	10.47
(%)		(0.5)	(13.7)	(33.7)	(33.2)	(13.7)	(4.4)	(1.0)					(100.0)				
腎形成不全		5	57	95	99	71	24	8	4	1			364	4	368	53.06	13.19
(%)		(1.4)	(15.7)	(26.1)	(27.2)	(19.5)	(6.6)	(2.2)	(1.1)	(0.3)			(100.0)				
不明		132	2,425	5,862	6,114	3,282	1,216	388	158	50	21	20	19,668	810	20,478	53.39	12.79
(%)		(0.7)	(12.3)	(29.8)	(31.1)	(16.7)	(6.2)	(2.0)	(0.8)	(0.3)	(0.1)	(0.1)	(100.0)				
再導入		5	98	330	383	198	78	19	11	5	1	1	1,129	35	1,164	54.61	12.47
(%)		(0.4)	(8.7)	(29.2)	(33.9)	(17.5)	(6.9)	(1.7)	(1.0)	(0.4)	(0.1)	(0.1)	(100.0)				
その他		37	645	1,581	1,602	897	305	111	35	21	7	8	5,249	157	5,406	53.45	12.98
(%)		(0.7)	(12.3)	(30.1)	(30.5)	(17.1)	(5.8)	(2.1)	(0.7)	(0.4)	(0.1)	(0.2)	(100.0)				
合計		761	17,496	53,890	64,737	40,392	16,459	6,163	2,216	844	303	192	203,453	6,083	209,536	55.77	13.19
(%)		(0.4)	(8.6)	(26.5)	(31.8)	(19.9)	(8.1)	(3.0)	(1.1)	(0.4)	(0.1)	(0.1)	(100.0)				
記載なし																	
総計		761	17,496	53,890	64,737	40,392	16,459	6,163	2,216	844	303	192	203,453	6,083	209,536	55.77	13.19
(%)		(0.4)	(8.6)	(26.5)	(31.8)	(19.9)	(8.1)	(3.0)	(1.1)	(0.4)	(0.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D2113##

(C)Japanese Society for Dialysis Therapy

2016年調査