

表 918 透析後体重(kg) 透析歴(年)別 (血液透析濾過患者 透析回数週3回)

透析歴	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
2年未満		16	547	2,068	3,085	2,220	1,082	471	171	84	26	20	9,790	228	10,018	59.02	14.24
(%)		(0.2)	(5.6)	(21.1)	(31.5)	(22.7)	(11.1)	(4.8)	(1.7)	(0.9)	(0.3)	(0.2)	(100.0)				
2年～		29	809	3,202	4,689	3,669	1,866	803	342	113	49	29	15,600	338	15,938	59.70	14.35
(%)		(0.2)	(5.2)	(20.5)	(30.1)	(23.5)	(12.0)	(5.1)	(2.2)	(0.7)	(0.3)	(0.2)	(100.0)				
5年～		33	988	3,737	5,296	4,054	1,859	742	282	109	40	27	17,167	415	17,582	58.59	13.80
(%)		(0.2)	(5.8)	(21.8)	(30.8)	(23.6)	(10.8)	(4.3)	(1.6)	(0.6)	(0.2)	(0.2)	(100.0)				
10年～		21	654	2,457	3,188	2,199	880	298	119	32	12	6	9,866	250	10,116	56.74	12.76
(%)		(0.2)	(6.6)	(24.9)	(32.3)	(22.3)	(8.9)	(3.0)	(1.2)	(0.3)	(0.1)	(0.1)	(100.0)				
15年～		13	465	1,720	2,011	1,193	380	108	30	5	5	1	5,931	127	6,058	54.60	11.41
(%)		(0.2)	(7.8)	(29.0)	(33.9)	(20.1)	(6.4)	(1.8)	(0.5)	(0.1)	(0.1)	(0.0)	(100.0)				
20年～		8	349	1,284	1,244	643	188	59	12	4	1	1	3,793	64	3,857	52.98	11.05
(%)		(0.2)	(9.2)	(33.9)	(32.8)	(17.0)	(5.0)	(1.6)	(0.3)	(0.1)	(0.0)	(0.0)	(100.0)				
25年～		6	249	818	792	322	89	13	3	2			2,294	51	2,345	51.61	10.06
(%)		(0.3)	(10.9)	(35.7)	(34.5)	(14.0)	(3.9)	(0.6)	(0.1)	(0.1)			(100.0)				
30年～		3	158	518	428	167	52	9					1,335	32	1,367	50.78	9.92
(%)		(0.2)	(11.8)	(38.8)	(32.1)	(12.5)	(3.9)	(0.7)					(100.0)				
35年～		1	121	282	252	86	21	2	1				766	14	780	49.84	9.61
(%)		(0.1)	(15.8)	(36.8)	(32.9)	(11.2)	(2.7)	(0.3)	(0.1)				(100.0)				
40年～		1	49	118	85	21	3	1					278	9	287	48.20	8.70
(%)		(0.4)	(17.6)	(42.4)	(30.6)	(7.6)	(1.1)	(0.4)					(100.0)				
合計		131	4,389	16,204	21,070	14,574	6,420	2,506	960	349	133	84	66,820	1,528	68,348	57.43	13.53
(%)		(0.2)	(6.6)	(24.3)	(31.5)	(21.8)	(9.6)	(3.8)	(1.4)	(0.5)	(0.2)	(0.1)	(100.0)				
不明			4	4	5	5	3						21	5	26	56.06	12.55
(%)			(19.0)	(19.0)	(23.8)	(23.8)	(14.3)						(100.0)				
記載なし																	
(%)																	
総計		131	4,393	16,208	21,075	14,579	6,423	2,506	960	349	133	84	66,841	1,533	68,374	57.43	13.53
(%)		(0.2)	(6.6)	(24.2)	(31.5)	(21.8)	(9.6)	(3.7)	(1.4)	(0.5)	(0.2)	(0.1)	(100.0)				
平均		9.89	11.64	10.88	9.38	8.05	6.91	5.89	5.42	4.83	4.94	4.57	9.14	8.96	9.14		
標準偏差		8.85	10.12	9.49	8.57	7.34	6.47	5.44	4.69	4.37	4.16	4.06	8.48	8.36	8.47		

数値下のかっこ内は行方向の合計に対する%です。
##D2122##
(C)Japanese Society for Dialysis Therapy
2016年調査