

表 924 体重(kg) 透析歴(年)別 (腹膜透析患者)

透析歴	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
2年未満	17	5	64	365	598	645	420	146	56	15	5	4	2,340	1,278	3,618	62.07	14.50
(%)	(0.7)	(0.2)	(2.7)	(15.6)	(25.6)	(27.6)	(17.9)	(6.2)	(2.4)	(0.6)	(0.2)	(0.2)	(100.0)				
2年～	20	5	46	240	569	607	286	103	27	9	7	2	1,921	1,063	2,984	61.49	13.61
(%)	(1.0)	(0.3)	(2.4)	(12.5)	(29.6)	(31.6)	(14.9)	(5.4)	(1.4)	(0.5)	(0.4)	(0.1)	(100.0)				
5年～	8	6	32	172	322	302	125	41	15	5	2	1	1,031	544	1,575	59.71	13.33
(%)	(0.8)	(0.6)	(3.1)	(16.7)	(31.2)	(29.3)	(12.1)	(4.0)	(1.5)	(0.5)	(0.2)	(0.1)	(100.0)				
10年～	2	3	8	42	75	49	20	9	1	1	1		211	78	289	57.55	13.60
(%)	(0.9)	(1.4)	(3.8)	(19.9)	(35.5)	(23.2)	(9.5)	(4.3)	(0.5)	(0.5)	(0.5)		(100.0)				
15年～			3	13	24	15	3						58	27	85	55.15	8.92
(%)			(5.2)	(22.4)	(41.4)	(25.9)	(5.2)						(100.0)				
20年～		1	3	6	4	4	1						19	9	28	50.85	13.58
(%)		(5.3)	(15.8)	(31.6)	(21.1)	(21.1)	(5.3)						(100.0)				
25年～				1	1	2							4	5	9	57.08	10.43
(%)				(25.0)	(25.0)	(50.0)							(100.0)				
30年～			1	2	2								5	1	6	44.88	9.33
(%)			(20.0)	(40.0)	(40.0)								(100.0)				
35年～														1	1		
(%)																	
40年～			1	1									2		2	39.70	4.24
(%)			(50.0)	(50.0)									(100.0)				
合計	47	20	158	842	1,595	1,624	855	299	99	30	15	7	5,591	3,006	8,597	61.13	13.97
(%)	(0.8)	(0.4)	(2.8)	(15.1)	(28.5)	(29.0)	(15.3)	(5.3)	(1.8)	(0.5)	(0.3)	(0.1)	(100.0)				
不明					2	1	2						5	13	18	65.40	9.35
(%)					(40.0)	(20.0)	(40.0)						(100.0)				
記載なし																	
(%)																	
総計	47	20	158	842	1,597	1,625	857	299	99	30	15	7	5,596	3,019	8,615	61.13	13.97
(%)	(0.8)	(0.4)	(2.8)	(15.0)	(28.5)	(29.0)	(15.3)	(5.3)	(1.8)	(0.5)	(0.3)	(0.1)	(100.0)				
平均	3.11	5.60	4.00	3.47	3.37	3.00	2.49	2.40	2.05	2.37	2.87	1.71	3.08	2.93	3.03		
標準偏差	3.11	5.52	5.70	4.38	3.74	3.25	2.77	2.64	2.28	2.76	2.83	2.21	3.60	3.36	3.52		

数値下のかっこ内は行方向の合計に対する%です。
##D2213##
(C)Japanese Society for Dialysis Therapy
2016年調査