

表 930 体重減少量(kg) 医療機関別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

医療機関	0.0未満	0.0~	1.0~	2.0~	3.0~	4.0~	5.0~	6.0~	7.0~	8.0~	合計	記載なし	総計	平均	標準偏差
国公立大学	4	40	112	93	38	2	1				290	4	294	1.88	0.95
(%)	(1.4)	(13.8)	(38.6)	(32.1)	(13.1)	(0.7)	(0.3)				(100.0)				
私立大学	2	69	265	356	190	32	6	3	1	924	106	1,030	2.32	1.01	
(%)	(0.2)	(7.5)	(28.7)	(38.5)	(20.6)	(3.5)	(0.6)	(0.3)	(0.1)	(100.0)					
国立病院機構等	2	15	84	112	66	13	3				295	41	336	2.43	1.00
(%)	(0.7)	(5.1)	(28.5)	(38.0)	(22.4)	(4.4)	(1.0)				(100.0)				
県市町村立・国保	37	425	2,193	3,847	1,821	354	45	6	2	8,730	681	9,411	2.40	0.92	
(%)	(0.4)	(4.9)	(25.1)	(44.1)	(20.9)	(4.1)	(0.5)	(0.1)	(0.0)	(100.0)					
JCHO(旧社会保険)等	13	83	435	808	400	84	16	3	2	1	1,845	22	1,867	2.45	0.97
(%)	(0.7)	(4.5)	(23.6)	(43.8)	(21.7)	(4.6)	(0.9)	(0.2)	(0.1)	(0.1)	(100.0)				
厚生連等	24	194	934	2,035	1,024	218	51	11	5	2	4,498	315	4,813	2.52	0.99
(%)	(0.5)	(4.3)	(20.8)	(45.2)	(22.8)	(4.8)	(1.1)	(0.2)	(0.1)	(0.0)	(100.0)				
その他公的	29	240	1,187	2,085	957	179	29	6	2		4,714	233	4,947	2.38	0.93
(%)	(0.6)	(5.1)	(25.2)	(44.2)	(20.3)	(3.8)	(0.6)	(0.1)	(0.0)		(100.0)				
私立総合	19	170	861	1,505	706	171	37	7	1	3	3,480	78	3,558	2.44	1.00
(%)	(0.5)	(4.9)	(24.7)	(43.2)	(20.3)	(4.9)	(1.1)	(0.2)	(0.0)	(0.1)	(100.0)				
私立	287	3,187	12,768	18,296	8,745	2,238	438	85	20	15	46,079	1,341	47,420	2.34	1.03
(%)	(0.6)	(6.9)	(27.7)	(39.7)	(19.0)	(4.9)	(1.0)	(0.2)	(0.0)	(0.0)	(100.0)				
私立診療所	322	3,420	20,037	36,686	19,075	5,015	1,026	229	55	12	85,877	1,580	87,457	2.51	0.99
(%)	(0.4)	(4.0)	(23.3)	(42.7)	(22.2)	(5.8)	(1.2)	(0.3)	(0.1)	(0.0)	(100.0)				
総計	739	7,843	38,876	65,823	33,022	8,306	1,652	350	87	34	156,732	4,401	161,133	2.45	1.00
(%)	(0.5)	(5.0)	(24.8)	(42.0)	(21.1)	(5.3)	(1.1)	(0.2)	(0.1)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D2314##

(C)Japanese Society for Dialysis Therapy

2016年調査