

表 935 体重減少量(kg) 医療機関別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

医療機関	0.0未満	0.0~	1.0~	2.0~	3.0~	4.0~	5.0~	6.0~	7.0~	8.0~	合計	記載なし	総計	平均	標準偏差
国公立大学	1	2	18	17	3	2					43	2	45	2.00	0.93
(%)	(2.3)	(4.7)	(41.9)	(39.5)	(7.0)	(4.7)					(100.0)				
私立大学		10	41	82	65	8		1			207	5	212	2.58	0.94
(%)		(4.8)	(19.8)	(39.6)	(31.4)	(3.9)		(0.5)			(100.0)				
国立病院機構等		1	12	13	6	1					33	13	46	2.38	0.93
(%)		(3.0)	(36.4)	(39.4)	(18.2)	(3.0)					(100.0)				
県市町村立・国保	9	81	472	933	501	96	23	1	1		2,117	95	2,212	2.49	0.93
(%)	(0.4)	(3.8)	(22.3)	(44.1)	(23.7)	(4.5)	(1.1)	(0.0)	(0.0)		(100.0)				
JCHO(旧社会保険)等	3	15	82	246	98	35	2	1		1	483	6	489	2.59	0.95
(%)	(0.6)	(3.1)	(17.0)	(50.9)	(20.3)	(7.2)	(0.4)	(0.2)		(0.2)	(100.0)				
厚生連等	4	41	226	563	306	67	11	5			1,223	68	1,291	2.61	0.94
(%)	(0.3)	(3.4)	(18.5)	(46.0)	(25.0)	(5.5)	(0.9)	(0.4)			(100.0)				
その他公的	1	32	211	395	206	51	15	4	1		916	65	981	2.56	1.00
(%)	(0.1)	(3.5)	(23.0)	(43.1)	(22.5)	(5.6)	(1.6)	(0.4)	(0.1)		(100.0)				
私立総合	2	49	282	528	252	63	14	2			1,192	45	1,237	2.48	0.95
(%)	(0.2)	(4.1)	(23.7)	(44.3)	(21.1)	(5.3)	(1.2)	(0.2)			(100.0)				
私立	76	714	3,654	6,150	3,375	812	172	31	10	3	14,997	347	15,344	2.47	1.01
(%)	(0.5)	(4.8)	(24.4)	(41.0)	(22.5)	(5.4)	(1.1)	(0.2)	(0.1)	(0.0)	(100.0)				
私立診療所	112	1,133	7,928	15,584	8,243	2,289	420	86	18	6	35,819	654	36,473	2.57	0.97
(%)	(0.3)	(3.2)	(22.1)	(43.5)	(23.0)	(6.4)	(1.2)	(0.2)	(0.1)	(0.0)	(100.0)				
総計	208	2,078	12,926	24,511	13,055	3,424	657	131	30	10	57,030	1,300	58,330	2.54	0.98
(%)	(0.4)	(3.6)	(22.7)	(43.0)	(22.9)	(6.0)	(1.2)	(0.2)	(0.1)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D2324##

(C)Japanese Society for Dialysis Therapy

2016年調査