

表 939 体重減少率(%) 原疾患別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

原疾患	1.0未満	1.0~	2.0~	3.0~	4.0~	5.0~	6.0~	7.0~	8.0~	9.0~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	862	1,611	4,480	9,310	12,890	10,497	5,359	1,978	667	294	47,948	1,285	49,233	4.55	1.62
(%)	(1.8)	(3.4)	(9.3)	(19.4)	(26.9)	(21.9)	(11.2)	(4.1)	(1.4)	(0.6)	(100.0)				
慢性腎盂腎炎	41	67	158	305	381	333	145	67	24	8	1,529	24	1,553	4.42	1.70
(%)	(2.7)	(4.4)	(10.3)	(19.9)	(24.9)	(21.8)	(9.5)	(4.4)	(1.6)	(0.5)	(100.0)				
急速進行性糸球体腎炎	30	45	123	211	293	256	137	52	14	8	1,169	42	1,211	4.51	1.73
(%)	(2.6)	(3.8)	(10.5)	(18.0)	(25.1)	(21.9)	(11.7)	(4.4)	(1.2)	(0.7)	(100.0)				
妊娠腎/妊娠中毒症	5	15	68	163	239	166	93	26	11	8	794	16	810	4.65	1.51
(%)	(0.6)	(1.9)	(8.6)	(20.5)	(30.1)	(20.9)	(11.7)	(3.3)	(1.4)	(1.0)	(100.0)				
その他の分類不能の腎炎	26	30	52	132	190	133	72	30	19	6	690	19	709	4.51	1.78
(%)	(3.8)	(4.3)	(7.5)	(19.1)	(27.5)	(19.3)	(10.4)	(4.3)	(2.8)	(0.9)	(100.0)				
多発性囊胞腎	113	247	742	1,447	1,713	1,148	479	153	38	16	6,096	141	6,237	4.23	1.49
(%)	(1.9)	(4.1)	(12.2)	(23.7)	(28.1)	(18.8)	(7.9)	(2.5)	(0.6)	(0.3)	(100.0)				
腎硬化症	317	626	1,536	2,991	3,803	2,940	1,344	541	171	80	14,349	342	14,691	4.38	1.63
(%)	(2.2)	(4.4)	(10.7)	(20.8)	(26.5)	(20.5)	(9.4)	(3.8)	(1.2)	(0.6)	(100.0)				
悪性高血圧	20	46	113	233	360	294	173	50	23	11	1,323	46	1,369	4.66	1.63
(%)	(1.5)	(3.5)	(8.5)	(17.6)	(27.2)	(22.2)	(13.1)	(3.8)	(1.7)	(0.8)	(100.0)				
糖尿病性腎症	1,248	2,319	6,420	12,469	16,584	12,430	5,794	2,185	699	350	60,498	1,739	62,237	4.42	1.61
(%)	(2.1)	(3.8)	(10.6)	(20.6)	(27.4)	(20.5)	(9.6)	(3.6)	(1.2)	(0.6)	(100.0)				
SLE腎炎	18	36	106	205	254	221	146	53	16	8	1,063	32	1,095	4.61	1.66
(%)	(1.7)	(3.4)	(10.0)	(19.3)	(23.9)	(20.8)	(13.7)	(5.0)	(1.5)	(0.8)	(100.0)				
アミロイド腎	10	10	19	28	51	43	21	7	3	192	4	196	4.35	1.84	
(%)	(5.2)	(5.2)	(9.9)	(14.6)	(26.6)	(22.4)	(10.9)	(3.6)	(1.6)	(100.0)					
痛風腎	16	26	53	124	162	113	58	22	2	2	578	19	597	4.33	1.63
(%)	(2.8)	(4.5)	(9.2)	(21.5)	(28.0)	(19.6)	(10.0)	(3.8)	(0.3)	(0.3)	(100.0)				
先天性代謝異常による腎不全	6	3	13	22	30	32	15	7	5	1	134	3	137	4.60	1.98
(%)	(4.5)	(2.2)	(9.7)	(16.4)	(22.4)	(23.9)	(11.2)	(5.2)	(3.7)	(0.7)	(100.0)				
腎・尿路結核	2	5	7	22	23	16	10	5			90	4	94	4.35	1.57
(%)	(2.2)	(5.6)	(7.8)	(24.4)	(25.6)	(17.8)	(11.1)	(5.6)			(100.0)				
腎・尿路結石	9	20	36	55	77	62	20	11	1	2	293	11	304	4.20	1.66
(%)	(3.1)	(6.8)	(12.3)	(18.8)	(26.3)	(21.2)	(6.8)	(3.8)	(0.3)	(0.7)	(100.0)				
腎・尿路腫瘍	12	22	41	80	123	94	37	19	5	2	435	7	442	4.41	1.64
(%)	(2.8)	(5.1)	(9.4)	(18.4)	(28.3)	(21.6)	(8.5)	(4.4)	(1.1)	(0.5)	(100.0)				
閉塞性尿路障害	21	12	44	55	84	63	39	20	6	3	347	8	355	4.39	1.93
(%)	(6.1)	(3.5)	(12.7)	(15.9)	(24.2)	(18.2)	(11.2)	(5.8)	(1.7)	(0.9)	(100.0)				
骨髄腫	9	6	11	14	21	17	9	3			90	2	92	3.86	2.15
(%)	(10.0)	(6.7)	(12.2)	(15.6)	(23.3)	(18.9)	(10.0)	(3.3)			(100.0)				
腎形成不全	9	12	23	60	78	71	40	23	3	3	322	4	326	4.69	1.75
(%)	(2.8)	(3.7)	(7.1)	(18.6)	(24.2)	(22.0)	(12.4)	(7.1)	(0.9)	(0.9)	(100.0)				
不明	393	599	1,518	2,816	3,637	2,849	1,409	604	212	105	14,142	534	14,676	4.42	1.74
(%)	(2.8)	(4.2)	(10.7)	(19.9)	(25.7)	(20.1)	(10.0)	(4.3)	(1.5)	(0.7)	(100.0)				
再導入	11	21	80	159	249	207	136	50	15	12	940	27	967	4.79	1.65
(%)	(1.2)	(2.2)	(8.5)	(16.9)	(26.5)	(22.0)	(14.5)	(5.3)	(1.6)	(1.3)	(100.0)				
その他	106	173	402	682	919	744	408	178	63	35	3,710	92	3,802	4.47	1.83
(%)	(2.9)	(4.7)	(10.8)	(18.4)	(24.8)	(20.1)	(11.0)	(4.8)	(1.7)	(0.9)	(100.0)				
合計	3,284	5,951	16,045	31,583	42,161	32,729	15,944	6,084	1,997	954	156,732	4,401	161,133	4.45	1.63
(%)	(2.1)	(3.8)	(10.2)	(20.2)	(26.9)	(20.9)	(10.2)	(3.9)	(1.3)	(0.6)	(100.0)				
記載なし															
(%)															
総計	3,284	5,951	16,045	31,583	42,161	32,729	15,944	6,084	1,997	954	156,732	4,401	161,133	4.45	1.63
(%)	(2.1)	(3.8)	(10.2)	(20.2)	(26.9)	(20.9)	(10.2)	(3.9)	(1.3)	(0.6)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D2413##

(C)Japanese Society for Dialysis Therapy

2016年調査