

表 946 Body Mass Index 治療方法別(透析患者全体)

治療方法	14未満	14~	16~	18~	18.5~	20~	22~	24~	25~	26~	28~	30~	35~	合計	記載なし	総計	平均	標準偏差
施設血液透析	1,284	7,263	22,961	9,071	32,054	44,323	34,847	12,230	9,238	11,646	6,196	5,494	45	196,652	37,052	233,704	21.50	3.73
(%)	(0.7)	(3.7)	(11.7)	(4.6)	(16.3)	(22.5)	(17.7)	(6.2)	(4.7)	(5.9)	(3.2)	(2.8)	(0.0)	(100.0)				
血液透析濾過	254	1,774	6,642	2,802	9,925	14,547	11,980	4,308	3,241	4,246	2,333	2,146	15	64,213	10,586	74,799	21.84	3.75
(%)	(0.4)	(2.8)	(10.3)	(4.4)	(15.5)	(22.7)	(18.7)	(6.7)	(5.0)	(6.6)	(3.6)	(3.3)	(0.0)	(100.0)				
血液濾過		3			3	2		1					9	10	19	20.90	3.22	
(%)		(33.3)			(33.3)	(22.2)		(11.1)					(100.0)					
血液吸着透析	8	43	221	77	264	323	172	57	37	31	11	11		1,255	100	1,355	20.31	3.03
(%)	(0.6)	(3.4)	(17.6)	(6.1)	(21.0)	(25.7)	(13.7)	(4.5)	(2.9)	(2.5)	(0.9)	(0.9)	(0.9)	(100.0)				
在宅血液透析	3	26	11	52	83	80	33	26	35	13	12		374	243	617	22.48	3.41	
(%)	(0.8)	(7.0)	(2.9)	(13.9)	(22.2)	(21.4)	(8.8)	(7.0)	(9.4)	(3.5)	(3.2)	(3.2)	(3.2)	(100.0)				
腹膜透析	30	62	217	115	498	1,106	1,211	562	432	560	298	241	4	5,336	3,279	8,615	23.22	3.65
(%)	(0.6)	(1.2)	(4.1)	(2.2)	(9.3)	(20.7)	(22.7)	(10.5)	(8.1)	(10.5)	(5.6)	(4.5)	(0.1)	(100.0)				
合計	1,576	9,145	30,070	12,076	42,793	60,385	48,292	17,190	12,975	16,518	8,851	7,904	64	267,839	51,270	319,109	21.61	3.74
(%)	(0.6)	(3.4)	(11.2)	(4.5)	(16.0)	(22.5)	(18.0)	(6.4)	(4.8)	(6.2)	(3.3)	(3.0)	(0.0)	(100.0)				
記載なし																		
(%)																		
総計	1,576	9,145	30,070	12,076	42,793	60,385	48,292	17,190	12,975	16,518	8,851	7,904	64	267,839	51,270	319,109	21.61	3.74
(%)	(0.6)	(3.4)	(11.2)	(4.5)	(16.0)	(22.5)	(18.0)	(6.4)	(4.8)	(6.2)	(3.3)	(3.0)	(0.0)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D2510##

(C)Japanese Society for Dialysis Therapy

2016年調査