

表 947 Body Mass Index 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	14未満	14~	16~	18~	18.5~	20~	22~	24~	25~	26~	28~	30~	35~	合計	記載なし	総計	平均	標準偏差
施設血液透析	1,284	7,263	22,961	9,071	32,054	44,323	34,847	12,230	9,238	11,646	6,196	5,494	45	196,652	37,052	233,704	21.50	3.73
(%)	(0.7)	(3.7)	(11.7)	(4.6)	(16.3)	(22.5)	(17.7)	(6.2)	(4.7)	(5.9)	(3.2)	(2.8)	(0.0)	(100.0)				
Off-line HDF	30	159	560	205	702	913	655	228	164	219	107	95		4,037	600	4,637	21.16	3.66
(%)	(0.7)	(3.9)	(13.9)	(5.1)	(17.4)	(22.6)	(16.2)	(5.6)	(4.1)	(5.4)	(2.7)	(2.4)		(100.0)				
On-line HDF	163	1,306	5,011	2,141	7,789	11,456	9,464	3,431	2,596	3,400	1,944	1,805	13	50,519	8,597	59,116	21.94	3.77
(%)	(0.3)	(2.6)	(9.9)	(4.2)	(15.4)	(22.7)	(18.7)	(6.8)	(5.1)	(6.7)	(3.8)	(3.6)	(0.0)	(100.0)				
Push/Pull HDF		3	9	5	11	12	11	3	3	2		1		60	6	66	20.71	3.40
(%)		(5.0)	(15.0)	(8.3)	(18.3)	(20.0)	(18.3)	(5.0)	(5.0)	(3.3)		(1.7)		(100.0)				
ハイオフィルトレーション	1	6	27	16	37	48	35	15	10	17	2	1		215	37	252	21.06	3.26
(%)	(0.5)	(2.8)	(12.6)	(7.4)	(17.2)	(22.3)	(16.3)	(7.0)	(4.7)	(7.9)	(0.9)	(0.5)		(100.0)				
IHDF	60	300	1,035	435	1,386	2,118	1,815	631	468	608	280	244	2	9,382	1,346	10,728	21.64	3.66
(%)	(0.6)	(3.2)	(11.0)	(4.6)	(14.8)	(22.6)	(19.3)	(6.7)	(5.0)	(6.5)	(3.0)	(2.6)	(0.0)	(100.0)				
血液濾過		3			3	2		1						9	10	19	20.90	3.22
(%)		(33.3)			(33.3)	(22.2)		(11.1)						(100.0)				
血液吸着透析 (リクセル等使用)	8	43	221	77	264	323	172	57	37	31	11	11		1,255	100	1,355	20.31	3.03
(%)	(0.6)	(3.4)	(17.6)	(6.1)	(21.0)	(25.7)	(13.7)	(4.5)	(2.9)	(2.5)	(0.9)	(0.9)		(100.0)				
在宅血液透析		3	26	11	52	83	80	33	26	35	13	12		374	243	617	22.48	3.41
(%)		(0.8)	(7.0)	(2.9)	(13.9)	(22.2)	(21.4)	(8.8)	(7.0)	(9.4)	(3.5)	(3.2)		(100.0)				
腹膜透析 (CAPD)	8	23	117	66	308	634	727	293	237	336	167	132	1	3,049	1,924	4,973	23.25	3.55
(%)	(0.3)	(0.8)	(3.8)	(2.2)	(10.1)	(20.8)	(23.8)	(9.6)	(7.8)	(11.0)	(5.5)	(4.3)	(0.0)	(100.0)				
腹膜透析 (APD)	18	29	84	35	134	350	327	185	133	165	96	72	2	1,630	1,007	2,637	23.09	3.84
(%)	(1.1)	(1.8)	(5.2)	(2.1)	(8.2)	(21.5)	(20.1)	(11.3)	(8.2)	(10.1)	(5.9)	(4.4)	(0.1)	(100.0)				
腹膜透析 (CCPD)	4	10	16	14	56	122	157	84	62	59	35	37	1	657	348	1,005	23.42	3.66
(%)	(0.6)	(1.5)	(2.4)	(2.1)	(8.5)	(18.6)	(23.9)	(12.8)	(9.4)	(9.0)	(5.3)	(5.6)	(0.2)	(100.0)				
合計	1,576	9,145	30,070	12,076	42,793	60,385	48,292	17,190	12,975	16,518	8,851	7,904	64	267,839	51,270	319,109	21.61	3.74
(%)	(0.6)	(3.4)	(11.2)	(4.5)	(16.0)	(22.5)	(18.0)	(6.4)	(4.8)	(6.2)	(3.3)	(3.0)	(0.0)	(100.0)				
記載なし																		
(%)																		
総計	1,576	9,145	30,070	12,076	42,793	60,385	48,292	17,190	12,975	16,518	8,851	7,904	64	267,839	51,270	319,109	21.61	3.74
(%)	(0.6)	(3.4)	(11.2)	(4.5)	(16.0)	(22.5)	(18.0)	(6.4)	(4.8)	(6.2)	(3.3)	(3.0)	(0.0)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

\*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

#D2511##

(C)Japanese Society for Dialysis Therapy

2016年調査