

表 983 BUN(mg/dL) 医療機関別 (腹膜透析患者)

医療機関	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
国公立大学		3	28	55	97	71	31	18	7	1		2	313	243	556	58.23	15.24
(%)		(1.0)	(8.9)	(17.6)	(31.0)	(22.7)	(9.9)	(5.8)	(2.2)	(0.3)		(0.6)	(100.0)				
私立大学	2	14	70	149	192	148	79	39	17	1	1	2	714	435	1,149	57.14	15.56
(%)	(0.3)	(2.0)	(9.8)	(20.9)	(26.9)	(20.7)	(11.1)	(5.5)	(2.4)	(0.1)	(0.1)	(0.3)	(100.0)				
国立病院機構等	1	6	18	45	51	36	19	7	2				185	42	227	54.14	14.30
(%)	(0.5)	(3.2)	(9.7)	(24.3)	(27.6)	(19.5)	(10.3)	(3.8)	(1.1)				(100.0)				
県市町村立・国保	4	30	132	268	334	212	101	38	9	2	1	1	1,132	468	1,600	54.24	14.63
(%)	(0.4)	(2.7)	(11.7)	(23.7)	(29.5)	(18.7)	(8.9)	(3.4)	(0.8)	(0.2)	(0.1)	(0.1)	(100.0)				
JCHO(旧社会保険)等	1	2	23	34	35	26	15	9	2	2			149	347	496	54.93	16.23
(%)	(0.7)	(1.3)	(15.4)	(22.8)	(23.5)	(17.4)	(10.1)	(6.0)	(1.3)	(1.3)			(100.0)				
厚生連等	1	10	52	90	95	60	39	15	5	3	1	1	372	179	551	55.01	18.26
(%)	(0.3)	(2.7)	(14.0)	(24.2)	(25.5)	(16.1)	(10.5)	(4.0)	(1.3)	(0.8)	(0.3)	(0.3)	(100.0)				
その他公的	3	37	121	214	246	175	87	27	9	4			923	404	1,327	53.66	14.78
(%)	(0.3)	(4.0)	(13.1)	(23.2)	(26.7)	(19.0)	(9.4)	(2.9)	(1.0)	(0.4)			(100.0)				
私立総合		12	42	70	93	79	37	16	5	2		1	357	36	393	55.68	16.25
(%)		(3.4)	(11.8)	(19.6)	(26.1)	(22.1)	(10.4)	(4.5)	(1.4)	(0.6)		(0.3)	(100.0)				
私立	5	25	145	251	261	199	107	57	14	8	3		1,075	551	1,626	54.98	16.02
(%)	(0.5)	(2.3)	(13.5)	(23.3)	(24.3)	(18.5)	(10.0)	(5.3)	(1.3)	(0.7)	(0.3)		(100.0)				
私立診療所	1	17	58	125	156	111	61	27	6	3			565	125	690	55.65	14.85
(%)	(0.2)	(3.0)	(10.3)	(22.1)	(27.6)	(19.6)	(10.8)	(4.8)	(1.1)	(0.5)			(100.0)				
総計	18	156	689	1,301	1,560	1,117	576	253	76	26	6	7	5,785	2,830	8,615	55.15	15.51
(%)	(0.3)	(2.7)	(11.9)	(22.5)	(27.0)	(19.3)	(10.0)	(4.4)	(1.3)	(0.4)	(0.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D2634##

(C)Japanese Society for Dialysis Therapy

2016年調査