

表 1012 equilibrated Kt/V(eKt/V) 原疾患別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

原疾患	0.4未満	0.4~	0.6~	0.8~	1.0~	1.2~	1.4~	1.6~	1.8~	2.0~	合計	記載なし	総計	平均	標準偏差		
慢性糸球体腎炎 (%)	37 (0.1)	60 (0.1)	396 (0.9)	2,418 (5.5)	8,753 (19.9)	14,086 (32.0)	10,459 (23.8)	4,954 (11.3)	1,814 (4.1)	1,035 (2.4)	44,012 (100.0)	5,221	49,233	1.36	0.28		
慢性腎盂腎炎 (%)	3 (0.2)	14 (1.0)	85 (6.1)	258 (18.6)	394 (28.4)	353 (25.4)	175 (12.6)	62 (4.5)	45 (3.2)	1,389 (100.0)	164	1,553	1.38	0.29			
急速進行性糸球体腎炎 (%)	1 (0.1)	10 (0.9)	54 (4.9)	223 (20.3)	329 (30.0)	252 (23.0)	146 (13.3)	56 (5.1)	26 (2.4)	1,097 (100.0)	114	1,211	1.38	0.28			
妊娠腎/妊娠中毒症 (%)	1 (0.1)	4 (0.5)	13 (1.7)	72 (9.6)	188 (25.1)	261 (34.8)	134 (17.9)	50 (6.7)	26 (3.5)	749 (100.0)	61	810	1.48	0.26			
その他の分類不能の腎炎 (%)	2 (0.3)	10 (1.6)	26 (4.1)	135 (21.3)	172 (27.2)	156 (24.6)	89 (14.1)	28 (4.4)	15 (2.4)	633 (100.0)	76	709	1.38	0.28			
多発性囊胞腎 (%)	4 (0.1)	8 (0.1)	57 (1.0)	413 (7.3)	1,421 (25.2)	1,852 (32.9)	1,196 (21.2)	467 (8.3)	139 (2.5)	77 (1.4)	5,634 (100.0)	603	6,237	1.31	0.26		
腎硬化症 (%)	11 (0.1)	34 (0.3)	184 (1.4)	1,097 (8.3)	3,319 (25.2)	4,236 (32.2)	2,630 (20.0)	1,123 (8.5)	359 (2.7)	168 (1.3)	13,161 (100.0)	1,530	14,691	1.30	0.26		
悪性高血圧 (%)	3 (0.2)	3 (0.2)	25 (2.0)	105 (8.5)	307 (24.9)	364 (29.5)	268 (21.8)	107 (8.7)	33 (2.7)	17 (1.4)	(100.0)	137	1,369	1.30	0.27		
糖尿病性腎症 (%)	50 (0.1)	162 (0.3)	1,116 (2.0)	5,828 (10.7)	16,257 (29.7)	17,811 (32.6)	9,040 (32.6)	3,033 (16.5)	934 (5.5)	433 (1.7)	54,664 (0.8)	7,573	62,237	1.25	0.25		
SLE腎炎 (%)			12 (1.2)	44 (4.5)	139 (14.3)	250 (25.7)	259 (26.6)	163 (16.7)	68 (7.0)	39 (4.0)	(100.0)	974	121	1,095	1.44	0.31	
アミロイド腎 (%)	1 (0.6)	1 (0.6)	1 (0.6)	12 (7.2)	40 (24.0)	41 (24.6)	37 (22.2)	19 (11.4)	10 (6.0)	5 (3.0)	(100.0)	167	29	196	1.35	0.33	
痛風腎 (%)			3 (0.6)	41 (7.7)	146 (27.4)	178 (33.5)	101 (19.0)	44 (8.3)	7 (1.3)	12 (2.3)	(100.0)	532	65	597	1.31	0.27	
先天性代謝異常による腎不全 (%)			4 (3.3)	27 (22.0)	38 (30.9)	27 (22.0)	18 (14.6)	3 (14.6)	6 (2.4)	123 (4.9)	(100.0)	14	137	1.41	0.31		
腎・尿路結核 (%)	1 (1.2)	1 (1.2)	5 (6.0)	14 (16.7)	26 (31.0)	21 (25.0)	12 (14.3)	3 (14.3)	1 (3.6)	84 (1.2)	(100.0)	10	94	1.35	0.28		
腎・尿路結石 (%)			7 (2.5)	18 (6.5)	50 (18.0)	108 (38.8)	58 (20.9)	22 (7.9)	10 (3.6)	278 (1.8)	(100.0)	26	304	1.33	0.27		
腎・尿路腫瘍 (%)	1 (0.2)	7 (1.7)	35 (8.6)	110 (27.2)	131 (32.3)	79 (19.5)	23 (5.7)	14 (3.5)	5 (1.2)	405 (100.0)	37	442	1.29	0.27			
閉塞性尿路障害 (%)	1 (0.3)	2 (0.6)	36 (11.1)	77 (23.8)	88 (27.2)	59 (18.3)	40 (12.4)	10 (3.1)	10 (3.1)	323 (100.0)	32	355	1.33	0.30			
骨髄腫 (%)			1 (1.2)	3 (3.6)	4 (4.8)	24 (28.9)	27 (32.5)	11 (13.3)	11 (13.3)	2 (2.4)	(100.0)	83	9	92	1.28	0.27	
腎形成不全 (%)			3 (1.0)	15 (5.1)	42 (14.3)	89 (30.3)	74 (25.2)	45 (15.3)	13 (4.4)	294 (4.4)	(100.0)	32	326	1.42	0.31		
不明 (%)	14 (0.1)	24 (0.2)	199 (1.6)	985 (7.8)	2,983 (23.7)	4,023 (32.0)	2,635 (21.0)	1,138 (9.1)	379 (3.0)	193 (1.5)	(100.0)	12,573	2,103	14,676	1.32	0.27	
再導入 (%)			2 (0.2)	4 (0.5)	52 (5.9)	163 (18.5)	265 (30.1)	217 (24.7)	108 (12.3)	40 (4.5)	(100.0)	29	880	87	967	1.39	0.30
その他 (%)	3 (0.1)	10 (0.3)	48 (1.5)	234 (7.1)	732 (22.2)	1,017 (30.9)	744 (22.6)	300 (9.1)	135 (4.1)	68 (2.1)	(100.0)	3,291	511	3,802	1.34	0.29	
合計 (%)	125 (0.1)	313 (0.2)	2,106 (1.5)	11,524 (8.1)	35,292 (24.8)	45,713 (32.1)	28,937 (20.3)	12,171 (8.5)	4,169 (2.9)	2,228 (1.6)	142,578 (100.0)	18,555	161,133	1.31	0.27		
記載なし (%)																	
総計 (%)	125 (0.1)	313 (0.2)	2,106 (1.5)	11,524 (8.1)	35,292 (24.8)	45,713 (32.1)	28,937 (20.3)	12,171 (8.5)	4,169 (2.9)	2,228 (1.6)	142,578 (100.0)	18,555	161,133	1.31	0.27		

数値下のかっこ内は行方向の合計に対する%です。

##D2913##

(C)Japanese Society for Dialysis Therapy

2016年調査