

表 1079 透析前アルブミン濃度(g/dL) 併用の状況別（透析患者全体）

併用の状況	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	177	1,031	4,473	17,021	74,326	137,654	42,782	2,094	140	279,698	30,440	310,138	3.56	0.44
(%)	(0.1)	(0.4)	(1.6)	(6.1)	(26.6)	(49.2)	(15.3)	(0.7)	(0.1)	(100.0)				
PD単独	18	75	261	793	1,690	1,325	314	20	4	4,500	2,431	6,931	3.25	0.54
(%)	(0.4)	(1.7)	(5.8)	(17.6)	(37.6)	(29.4)	(7.0)	(0.4)	(0.1)	(100.0)				
HD(F)単独.PDカテあり	2	6	14	15	46	46	22	1		152	25	177	3.27	0.69
(%)	(1.3)	(3.9)	(9.2)	(9.9)	(30.3)	(30.3)	(14.5)	(0.7)		(100.0)				
PD.HD併用週1HD(F)	4	4	51	242	514	318	33			1,166	330	1,496	3.20	0.44
(%)	(0.3)	(0.3)	(4.4)	(20.8)	(44.1)	(27.3)	(2.8)			(100.0)				
PD.HD併用週2HD(F)		1	4	14	65	46	6	1		137	28	165	3.33	0.42
(%)		(0.7)	(2.9)	(10.2)	(47.4)	(33.6)	(4.4)	(0.7)		(100.0)				
PD.HD併用週3HD(F)		2	3	3	6	15	3			32	5	37	3.31	0.73
(%)		(6.3)	(9.4)	(9.4)	(18.8)	(46.9)	(9.4)			(100.0)				
PD.HD併用週4HD(F)						1				1		1	3.60	
(%)						(100.0)				(100.0)				
PD.HD併用その他			2	11	16	18	1			48	15	63	3.26	0.46
(%)			(4.2)	(22.9)	(33.3)	(37.5)	(2.1)			(100.0)				
合計	201	1,119	4,808	18,099	76,663	139,423	43,161	2,116	144	285,734	33,274	319,008	3.55	0.44
(%)	(0.1)	(0.4)	(1.7)	(6.3)	(26.8)	(48.8)	(15.1)	(0.7)	(0.1)	(100.0)				
不明			4	4	5	4				17	5	22	2.92	0.62
(%)			(23.5)	(23.5)	(29.4)	(23.5)				(100.0)				
記載なし											79	79		
(%)														
総計	201	1,119	4,812	18,103	76,668	139,427	43,161	2,116	144	285,751	33,358	319,109	3.55	0.44
(%)	(0.1)	(0.4)	(1.7)	(6.3)	(26.8)	(48.8)	(15.1)	(0.7)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D3812##

(C)Japanese Society for Dialysis Therapy

2016年調査