

表 1088 透析前アルブミン濃度(g/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	35	170	756	3,015	14,248	27,260	8,896	432	36	54,848	2,324	57,172	3.58	0.43
(%)	(0.1)	(0.3)	(1.4)	(5.5)	(26.0)	(49.7)	(16.2)	(0.8)	(0.1)	(100.0)				
慢性腎盂腎炎	1	6	35	125	503	874	255	11		1,810	58	1,868	3.53	0.44
(%)	(0.1)	(0.3)	(1.9)	(6.9)	(27.8)	(48.3)	(14.1)	(0.6)		(100.0)				
急速進行性糸球体腎炎		11	41	162	577	638	176	5	1	1,611	76	1,687	3.43	0.47
(%)		(0.7)	(2.5)	(10.1)	(35.8)	(39.6)	(10.9)	(0.3)	(0.1)	(100.0)				
妊娠腎/妊娠中毒症		1	8	41	202	425	127	7		811	38	849	3.60	0.40
(%)		(0.1)	(1.0)	(5.1)	(24.9)	(52.4)	(15.7)	(0.9)		(100.0)				
その他の分類不能の腎炎	1	1	12	56	207	413	157	19	1	867	35	902	3.61	0.46
(%)	(0.1)	(0.1)	(1.4)	(6.5)	(23.9)	(47.6)	(18.1)	(2.2)	(0.1)	(100.0)				
多発性嚢胞腎	4	20	91	391	2,026	3,647	1,045	53	3	7,280	268	7,548	3.56	0.41
(%)	(0.1)	(0.3)	(1.3)	(5.4)	(27.8)	(50.1)	(14.4)	(0.7)	(0.0)	(100.0)				
腎硬化症	15	78	348	1,624	6,427	9,620	2,455	111	7	20,685	833	21,518	3.50	0.43
(%)	(0.1)	(0.4)	(1.7)	(7.9)	(31.1)	(46.5)	(11.9)	(0.5)	(0.0)	(100.0)				
悪性高血圧		4	19	77	404	849	306	23	1	1,683	94	1,777	3.61	0.42
(%)		(0.2)	(1.1)	(4.6)	(24.0)	(50.4)	(18.2)	(1.4)	(0.1)	(100.0)				
糖尿病性腎症	54	374	1,592	5,475	21,300	39,165	12,554	571	40	81,125	3,343	84,468	3.55	0.45
(%)	(0.1)	(0.5)	(2.0)	(6.7)	(26.3)	(48.3)	(15.5)	(0.7)	(0.0)	(100.0)				
SLE腎炎	2	2	22	123	425	586	142	8		1,310	60	1,370	3.47	0.44
(%)	(0.2)	(0.2)	(1.7)	(9.4)	(32.4)	(44.7)	(10.8)	(0.6)		(100.0)				
アミロイド腎		5	11	45	85	94	22	7		269	11	280	3.34	0.55
(%)		(1.9)	(4.1)	(16.7)	(31.6)	(34.9)	(8.2)	(2.6)		(100.0)				
痛風腎			10	28	159	355	110	7		669	31	700	3.61	0.40
(%)			(1.5)	(4.2)	(23.8)	(53.1)	(16.4)	(1.0)		(100.0)				
先天性代謝異常による腎不全		3	4	10	34	70	41	3		165	4	169	3.60	0.57
(%)		(1.8)	(2.4)	(6.1)	(20.6)	(42.4)	(24.8)	(1.8)		(100.0)				
腎・尿路結核		1	3	10	28	47	11			100	6	106	3.44	0.47
(%)		(1.0)	(3.0)	(10.0)	(28.0)	(47.0)	(11.0)			(100.0)				
腎・尿路結石		2	7	23	120	169	45	1	1	368	16	384	3.51	0.45
(%)		(0.5)	(1.9)	(6.3)	(32.6)	(45.9)	(12.2)	(0.3)	(0.3)	(100.0)				
腎・尿路腫瘍		1	15	51	171	324	61	4	1	628	16	644	3.50	0.44
(%)		(0.2)	(2.4)	(8.1)	(27.2)	(51.6)	(9.7)	(0.6)	(0.2)	(100.0)				
閉塞性尿路障害		1	12	30	110	211	75	4		443	22	465	3.57	0.46
(%)		(0.2)	(2.7)	(6.8)	(24.8)	(47.6)	(16.9)	(0.9)		(100.0)				
骨髄腫	1	2	10	27	70	68	19	3		200	13	213	3.34	0.54
(%)	(0.5)	(1.0)	(5.0)	(13.5)	(35.0)	(34.0)	(9.5)	(1.5)		(100.0)				
腎形成不全		1	2	12	61	168	106	6	1	357	11	368	3.73	0.43
(%)		(0.3)	(0.6)	(3.4)	(17.1)	(47.1)	(29.7)	(1.7)	(0.3)	(100.0)				
不明	17	106	461	1,620	5,675	8,858	2,635	145	10	19,527	951	20,478	3.50	0.47
(%)	(0.1)	(0.5)	(2.4)	(8.3)	(29.1)	(45.4)	(13.5)	(0.7)	(0.1)	(100.0)				
再導入		2	10	48	247	579	219	13	1	1,119	45	1,164	3.64	0.41
(%)		(0.2)	(0.9)	(4.3)	(22.1)	(51.7)	(19.6)	(1.2)	(0.1)	(100.0)				
その他	11	39	141	410	1,447	2,310	802	53	5	5,218	188	5,406	3.51	0.50
(%)	(0.2)	(0.7)	(2.7)	(7.9)	(27.7)	(44.3)	(15.4)	(1.0)	(0.1)	(100.0)				
合計	141	830	3,610	13,403	54,526	96,730	30,259	1,486	108	201,093	8,443	209,536	3.54	0.45
(%)	(0.1)	(0.4)	(1.8)	(6.7)	(27.1)	(48.1)	(15.0)	(0.7)	(0.1)	(100.0)				
記載なし														
(%)														
総計	141	830	3,610	13,403	54,526	96,730	30,259	1,486	108	201,093	8,443	209,536	3.54	0.45
(%)	(0.1)	(0.4)	(1.8)	(6.7)	(27.1)	(48.1)	(15.0)	(0.7)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D3823##

(C)Japanese Society for Dialysis Therapy

2016年調査