

表 1094 透析前アルブミン濃度(g/dL) 医療機関別 (血液透析濾過患者 透析回数週3回)

医療機関	1.5未満	1.5~	2.0~	2.5~	3.0~	3.5~	4.0~	4.5~	5.0~	合計	記載なし	総計	平均	標準偏差
国公立大学		3	7	9	11	17	4			51	3	54	3.13	0.66
(%)		(5.9)	(13.7)	(17.6)	(21.6)	(33.3)	(7.8)			(100.0)				
私立大学	1	3	3	21	57	126	23			234	3	237	3.47	0.48
(%)	(0.4)	(1.3)	(1.3)	(9.0)	(24.4)	(53.8)	(9.8)			(100.0)				
国立病院・機構等	1	1	6	10	7	20	9	1		55		55	3.35	0.71
(%)	(1.8)	(1.8)	(10.9)	(18.2)	(12.7)	(36.4)	(16.4)	(1.8)		(100.0)				
県市町村立・国保		7	38	161	771	1,186	269	9	2	2,443	115	2,558	3.51	0.42
(%)		(0.3)	(1.6)	(6.6)	(31.6)	(48.5)	(11.0)	(0.4)	(0.1)	(100.0)				
JCHO(旧社会保険)等			8	30	164	248	64	4	1	519	7	526	3.54	0.40
(%)			(1.5)	(5.8)	(31.6)	(47.8)	(12.3)	(0.8)	(0.2)	(100.0)				
厚生連等		1	17	59	426	702	162	13	2	1,382	89	1,471	3.55	0.39
(%)		(0.1)	(1.2)	(4.3)	(30.8)	(50.8)	(11.7)	(0.9)	(0.1)	(100.0)				
その他公的		6	11	46	293	513	150	8	1	1,028	54	1,082	3.57	0.42
(%)		(0.6)	(1.1)	(4.5)	(28.5)	(49.9)	(14.6)	(0.8)	(0.1)	(100.0)				
私立総合	2	1	19	95	440	640	167	2	2	1,368	97	1,465	3.51	0.41
(%)	(0.1)	(0.1)	(1.4)	(6.9)	(32.2)	(46.8)	(12.2)	(0.1)	(0.1)	(100.0)				
私立	16	52	261	1,029	4,790	8,709	2,589	105	4	17,555	450	18,005	3.55	0.43
(%)	(0.1)	(0.3)	(1.5)	(5.9)	(27.3)	(49.6)	(14.7)	(0.6)	(0.0)	(100.0)				
私立診療所		20	150	1,273	9,759	23,297	6,994	324	13	41,830	1,091	42,921	3.64	0.36
(%)		(0.0)	(0.4)	(3.0)	(23.3)	(55.7)	(16.7)	(0.8)	(0.0)	(100.0)				
総計	20	94	520	2,733	16,718	35,458	10,431	466	25	66,465	1,909	68,374	3.60	0.39
(%)	(0.0)	(0.1)	(0.8)	(4.1)	(25.2)	(53.3)	(15.7)	(0.7)	(0.0)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

#D3834##

(C)Japanese Society for Dialysis Therapy

2016年調査