

表 1101 透析前CRP濃度(mg/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	0.2未満	0.2~	0.5~	1.0~	2.0~	4.0~	6.0~	8.0~	10.0~	15.0~	合計	記載なし	総計	平均	標準偏差
施設血液透析	97,901	36,044	18,660	12,363	7,338	2,690	1,389	704	815	518	178,422	55,282	233,704	0.67	1.90
(%)	(54.9)	(20.2)	(10.5)	(6.9)	(4.1)	(1.5)	(0.8)	(0.4)	(0.5)	(0.3)	(100.0)				
Off-line HDF	1,789	768	441	289	188	75	38	13	20	17	3,638	999	4,637	0.83	2.25
(%)	(49.2)	(21.1)	(12.1)	(7.9)	(5.2)	(2.1)	(1.0)	(0.4)	(0.5)	(0.5)	(100.0)				
On-line HDF	28,031	9,377	4,667	2,934	1,589	523	231	129	123	72	47,676	11,440	59,116	0.52	1.44
(%)	(58.8)	(19.7)	(9.8)	(6.2)	(3.3)	(1.1)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
Push/Pull HDF	18	10	5	2	4			1			40	26	66	0.77	1.57
(%)	(45.0)	(25.0)	(12.5)	(5.0)	(10.0)			(2.5)			(100.0)				
バイオフィルトレーション	102	45	24	19	10	8	1			2	211	41	252	0.88	2.37
(%)	(48.3)	(21.3)	(11.4)	(9.0)	(4.7)	(3.8)	(0.5)			(0.9)	(100.0)				
IHDF	4,620	1,804	978	618	379	113	40	23	22	13	8,610	2,118	10,728	0.59	1.49
(%)	(53.7)	(21.0)	(11.4)	(7.2)	(4.4)	(1.3)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
血液濾過	1	5	1	1	2		1			1	12	7	19	3.87	8.40
(%)	(8.3)	(41.7)	(8.3)	(8.3)	(16.7)		(8.3)			(8.3)	(100.0)				
血液吸着透析 (リクセル等使用)	574	253	136	80	51	15	6	1	4	4	1,124	231	1,355	0.63	1.55
(%)	(51.1)	(22.5)	(12.1)	(7.1)	(4.5)	(1.3)	(0.5)	(0.1)	(0.4)	(0.4)	(100.0)				
在宅血液透析	268	79	40	14	9	4	1		4	2	421	196	617	0.55	1.84
(%)	(63.7)	(18.8)	(9.5)	(3.3)	(2.1)	(1.0)	(0.2)		(1.0)	(0.5)	(100.0)				
腹膜透析 (CAPD)	1,620	557	319	224	113	47	21	10	16	6	2,933	2,040	4,973	0.66	1.64
(%)	(55.2)	(19.0)	(10.9)	(7.6)	(3.9)	(1.6)	(0.7)	(0.3)	(0.5)	(0.2)	(100.0)				
腹膜透析 (APD)	921	307	132	118	63	17	10	7	5	4	1,584	1,053	2,637	0.60	1.60
(%)	(58.1)	(19.4)	(8.3)	(7.4)	(4.0)	(1.1)	(0.6)	(0.4)	(0.3)	(0.3)	(100.0)				
腹膜透析 (CCPD)	377	111	74	37	21	2	3	2	2		629	376	1,005	0.48	1.16
(%)	(59.9)	(17.6)	(11.8)	(5.9)	(3.3)	(0.3)	(0.5)	(0.3)	(0.3)		(100.0)				
合計	136,222	49,360	25,477	16,699	9,767	3,494	1,741	890	1,011	639	245,300	73,809	319,109	0.64	1.80
(%)	(55.5)	(20.1)	(10.4)	(6.8)	(4.0)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし															
(%)															
総計	136,222	49,360	25,477	16,699	9,767	3,494	1,741	890	1,011	639	245,300	73,809	319,109	0.64	1.80
(%)	(55.5)	(20.1)	(10.4)	(6.8)	(4.0)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

##D3911##

(C)Japanese Society for Dialysis Therapy

2016年調査