

表 1102 透析前CRP濃度(mg/dL) 併用の状況別 (透析患者全体)

併用の状況	0.2未満	0.2~	0.5~	1.0~	2.0~	4.0~	6.0~	8.0~	10.0~	15.0~	合計	記載なし	総計	平均	標準偏差
HD(F)単独	133,193	48,336	24,928	16,303	9,552	3,422	1,705	870	986	627	239,922	70,216	310,138	0.64	1.81
(%)	(55.5)	(20.1)	(10.4)	(6.8)	(4.0)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
PD単独	2,277	753	406	282	157	51	26	13	16	9	3,990	2,941	6,931	0.61	1.59
(%)	(57.1)	(18.9)	(10.2)	(7.1)	(3.9)	(1.3)	(0.7)	(0.3)	(0.4)	(0.2)	(100.0)				
HD(F)単独.PDカテあり	61	35	12	11	9	4	1	1	1	2	137	40	177	1.16	2.96
(%)	(44.5)	(25.5)	(8.8)	(8.0)	(6.6)	(2.9)	(0.7)	(0.7)	(0.7)	(1.5)	(100.0)				
PD.HD併用週1HD(F)	581	204	106	85	32	14	6	5	7	1	1,041	455	1,496	0.63	1.55
(%)	(55.8)	(19.6)	(10.2)	(8.2)	(3.1)	(1.3)	(0.6)	(0.5)	(0.7)	(0.1)	(100.0)				
PD.HD併用週2HD(F)	61	16	17	11	9	2	1	1			118	47	165	0.73	1.37
(%)	(51.7)	(13.6)	(14.4)	(9.3)	(7.6)	(1.7)	(0.8)	(0.8)			(100.0)				
PD.HD併用週3HD(F)	16	4	3	2	4		1				30	7	37	0.86	1.46
(%)	(53.3)	(13.3)	(10.0)	(6.7)	(13.3)		(3.3)				(100.0)				
PD.HD併用週4HD(F)	1										1		1	0.05	
(%)	(100.0)										(100.0)				
PD.HD併用その他	24	10	3	4	2						43	20	63	0.42	0.67
(%)	(55.8)	(23.3)	(7.0)	(9.3)	(4.7)						(100.0)				
合計	136,214	49,358	25,475	16,698	9,765	3,493	1,740	890	1,010	639	245,282	73,726	319,008	0.64	1.80
(%)	(55.5)	(20.1)	(10.4)	(6.8)	(4.0)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
不明	8	2	2	1	2	1	1		1		18	4	22	1.90	3.21
(%)	(44.4)	(11.1)	(11.1)	(5.6)	(11.1)	(5.6)	(5.6)		(5.6)		(100.0)				
記載なし											79	79			
(%)															
総計	136,222	49,360	25,477	16,699	9,767	3,494	1,741	890	1,011	639	245,300	73,809	319,109	0.64	1.80
(%)	(55.5)	(20.1)	(10.4)	(6.8)	(4.0)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D3912##

(C)Japanese Society for Dialysis Therapy

2016年調査