

表 1107 透析前CRP濃度(mg/dL) 医療機関別 (透析患者全体)

医療機関	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	294	138	84	79	61	29	19	16	24	19	763	361	1,124	2.04	4.39
(%)	(38.5)	(18.1)	(11.0)	(10.4)	(8.0)	(3.8)	(2.5)	(2.1)	(3.1)	(2.5)	(100.0)				
私立大学	1,105	392	229	193	133	84	42	16	32	23	2,249	1,437	3,686	1.23	3.16
(%)	(49.1)	(17.4)	(10.2)	(8.6)	(5.9)	(3.7)	(1.9)	(0.7)	(1.4)	(1.0)	(100.0)				
国立病院機構等	241	95	49	46	38	13	8	5	5	4	504	407	911	1.11	2.55
(%)	(47.8)	(18.8)	(9.7)	(9.1)	(7.5)	(2.6)	(1.6)	(1.0)	(1.0)	(0.8)	(100.0)				
県市町村立・国保	7,799	2,800	1,420	993	601	250	139	82	93	51	14,228	5,520	19,748	0.75	2.21
(%)	(54.8)	(19.7)	(10.0)	(7.0)	(4.2)	(1.8)	(1.0)	(0.6)	(0.7)	(0.4)	(100.0)				
JCHO(旧社会保険)等	1,445	529	315	212	127	50	26	9	16	8	2,737	1,028	3,765	0.75	1.99
(%)	(52.8)	(19.3)	(11.5)	(7.7)	(4.6)	(1.8)	(0.9)	(0.3)	(0.6)	(0.3)	(100.0)				
厚生連等	4,125	1,445	733	497	289	101	68	28	33	22	7,341	2,044	9,385	0.66	1.77
(%)	(56.2)	(19.7)	(10.0)	(6.8)	(3.9)	(1.4)	(0.9)	(0.4)	(0.4)	(0.3)	(100.0)				
その他公的	3,961	1,414	745	506	308	130	75	31	56	42	7,268	3,590	10,858	0.80	2.28
(%)	(54.5)	(19.5)	(10.3)	(7.0)	(4.2)	(1.8)	(1.0)	(0.4)	(0.8)	(0.6)	(100.0)				
私立総合	3,360	1,165	635	400	272	89	51	22	28	26	6,048	1,765	7,813	0.71	2.33
(%)	(55.6)	(19.3)	(10.5)	(6.6)	(4.5)	(1.5)	(0.8)	(0.4)	(0.5)	(0.4)	(100.0)				
私立	37,850	14,788	8,087	5,617	3,482	1,355	671	343	415	261	72,869	18,720	91,589	0.76	2.03
(%)	(51.9)	(20.3)	(11.1)	(7.7)	(4.8)	(1.9)	(0.9)	(0.5)	(0.6)	(0.4)	(100.0)				
私立診療所	76,042	26,594	13,180	8,156	4,456	1,393	642	338	309	183	131,293	38,937	170,230	0.52	1.45
(%)	(57.9)	(20.3)	(10.0)	(6.2)	(3.4)	(1.1)	(0.5)	(0.3)	(0.2)	(0.1)	(100.0)				
総計	136,222	49,360	25,477	16,699	9,767	3,494	1,741	890	1,011	639	245,300	73,809	319,109	0.64	1.80
(%)	(55.5)	(20.1)	(10.4)	(6.8)	(4.0)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

##D3917##

(C)Japanese Society for Dialysis Therapy

2016年調査