

表 1111 透析前CRP濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	26,869	9,113	4,471	2,847	1,650	558	296	158	159	112	46,233	10,939	57,172	0.59	1.76
(%)	(58.1)	(19.7)	(9.7)	(6.2)	(3.6)	(1.2)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
慢性腎盂腎炎	858	298	170	95	55	27	18	3	10	1	1,535	333	1,868	0.64	1.61
(%)	(55.9)	(19.4)	(11.1)	(6.2)	(3.6)	(1.8)	(1.2)	(0.2)	(0.7)	(0.1)	(100.0)				
急速進行性糸球体腎炎	752	273	157	109	74	30	12	6	4	5	1,422	265	1,687	0.74	2.00
(%)	(52.9)	(19.2)	(11.0)	(7.7)	(5.2)	(2.1)	(0.8)	(0.4)	(0.3)	(0.4)	(100.0)				
妊娠腎/妊娠中毒症	408	134	45	40	25	8	2	2	2	1	667	182	849	0.53	1.79
(%)	(61.2)	(20.1)	(6.7)	(6.0)	(3.7)	(1.2)	(0.3)	(0.3)	(0.3)	(0.1)	(100.0)				
その他の分類不能の腎炎	453	135	71	46	27	3	1	4	1	3	744	158	902	0.52	1.59
(%)	(60.9)	(18.1)	(9.5)	(6.2)	(3.6)	(0.4)	(0.1)	(0.5)	(0.1)	(0.4)	(100.0)				
多発性嚢胞腎	3,322	1,301	683	421	273	90	59	22	29	28	6,228	1,320	7,548	0.74	2.26
(%)	(53.3)	(20.9)	(11.0)	(6.8)	(4.4)	(1.4)	(0.9)	(0.4)	(0.5)	(0.4)	(100.0)				
腎硬化症	9,401	3,686	1,981	1,355	744	279	132	59	76	46	17,759	3,759	21,518	0.68	1.95
(%)	(52.9)	(20.8)	(11.2)	(7.6)	(4.2)	(1.6)	(0.7)	(0.3)	(0.4)	(0.3)	(100.0)				
悪性高血圧	849	266	151	88	60	12	5	5	8	2	1,446	331	1,777	0.55	1.43
(%)	(58.7)	(18.4)	(10.4)	(6.1)	(4.1)	(0.8)	(0.3)	(0.3)	(0.6)	(0.1)	(100.0)				
糖尿病性腎症	37,495	14,141	7,333	4,958	2,894	1,085	561	271	335	180	69,253	15,215	84,468	0.67	1.81
(%)	(54.1)	(20.4)	(10.6)	(7.2)	(4.2)	(1.6)	(0.8)	(0.4)	(0.5)	(0.3)	(100.0)				
SLE腎炎	580	227	131	83	55	26	7	8	1	8	1,126	244	1,370	0.80	2.25
(%)	(51.5)	(20.2)	(11.6)	(7.4)	(4.9)	(2.3)	(0.6)	(0.7)	(0.1)	(0.7)	(100.0)				
アミロイド腎	125	51	23	21	16	5	2	3	3		249	31	280	0.86	1.87
(%)	(50.2)	(20.5)	(9.2)	(8.4)	(6.4)	(2.0)	(0.8)	(1.2)	(1.2)		(100.0)				
痛風腎	300	137	55	48	28	4	6	2	1	2	583	117	700	0.63	1.46
(%)	(51.5)	(23.5)	(9.4)	(8.2)	(4.8)	(0.7)	(1.0)	(0.3)	(0.2)	(0.3)	(100.0)				
先天性代謝異常による腎不全	81	20	11	10	3	3	2	1		3	134	35	169	1.23	4.49
(%)	(60.4)	(14.9)	(8.2)	(7.5)	(2.2)	(2.2)	(1.5)	(0.7)		(2.2)	(100.0)				
腎・尿路結核	49	11	9	4	5	2	2				82	24	106	0.68	1.38
(%)	(59.8)	(13.4)	(11.0)	(4.9)	(6.1)	(2.4)	(2.4)				(100.0)				
腎・尿路結石	157	60	36	24	17	7	2	2	1		306	78	384	0.70	1.43
(%)	(51.3)	(19.6)	(11.8)	(7.8)	(5.6)	(2.3)	(0.7)	(0.7)	(0.3)		(100.0)				
腎・尿路腫瘍	264	125	52	40	28	12	7	2	4	7	541	103	644	1.00	2.65
(%)	(48.8)	(23.1)	(9.6)	(7.4)	(5.2)	(2.2)	(1.3)	(0.4)	(0.7)	(1.3)	(100.0)				
閉塞性尿路障害	203	88	44	24	14	8	5		1	2	389	76	465	0.68	1.73
(%)	(52.2)	(22.6)	(11.3)	(6.2)	(3.6)	(2.1)	(1.3)		(0.3)	(0.5)	(100.0)				
骨髄腫	80	35	24	18	9	7	4	1		1	179	34	213	0.98	2.09
(%)	(44.7)	(19.6)	(13.4)	(10.1)	(5.0)	(3.9)	(2.2)	(0.6)		(0.6)	(100.0)				
腎形成不全	193	64	30	9	8	2	2	1			309	59	368	0.38	0.91
(%)	(62.5)	(20.7)	(9.7)	(2.9)	(2.6)	(0.6)	(0.6)	(0.3)			(100.0)				
不明	8,857	3,369	1,795	1,163	774	284	137	77	97	51	16,604	3,874	20,478	0.73	2.15
(%)	(53.3)	(20.3)	(10.8)	(7.0)	(4.7)	(1.7)	(0.8)	(0.5)	(0.6)	(0.3)	(100.0)				
再導入	525	186	108	77	33	11	6	4	6	4	960	204	1,164	0.66	1.79
(%)	(54.7)	(19.4)	(11.3)	(8.0)	(3.4)	(1.1)	(0.6)	(0.4)	(0.6)	(0.4)	(100.0)				
その他	2,318	903	509	354	199	81	41	33	28	22	4,488	918	5,406	0.81	2.15
(%)	(51.6)	(20.1)	(11.3)	(7.9)	(4.4)	(1.8)	(0.9)	(0.7)	(0.6)	(0.5)	(100.0)				
合計	94,139	34,623	17,889	11,834	6,991	2,544	1,309	664	766	478	171,237	38,299	209,536	0.66	1.88
(%)	(55.0)	(20.2)	(10.4)	(6.9)	(4.1)	(1.5)	(0.8)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし															
(%)															
総計	94,139	34,623	17,889	11,834	6,991	2,544	1,309	664	766	478	171,237	38,299	209,536	0.66	1.88
(%)	(55.0)	(20.2)	(10.4)	(6.9)	(4.1)	(1.5)	(0.8)	(0.4)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

##D3923##

(C)Japanese Society for Dialysis Therapy

2016年調査