

表 1116 透析前CRP濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	11,544	3,617	1,789	1,106	584	214	73	41	33	21	19,022	3,366	22,388	0.48	1.32
(%)	(60.7)	(19.0)	(9.4)	(5.8)	(3.1)	(1.1)	(0.4)	(0.2)	(0.2)	(0.1)	(100.0)				
慢性腎盂腎炎	326	103	46	34	17	2	3		4	3	538	89	627	0.60	1.91
(%)	(60.6)	(19.1)	(8.6)	(6.3)	(3.2)	(0.4)	(0.6)		(0.7)	(0.6)	(100.0)				
急速進行性糸球体腎炎	242	84	47	20	17	11	1	1	2		425	59	484	0.56	1.25
(%)	(56.9)	(19.8)	(11.1)	(4.7)	(4.0)	(2.6)	(0.2)	(0.2)	(0.5)		(100.0)				
妊娠腎/妊娠中毒症	247	73	29	27	12	3		1	1	2	395	64	459	0.52	1.64
(%)	(62.5)	(18.5)	(7.3)	(6.8)	(3.0)	(0.8)		(0.3)	(0.3)	(0.5)	(100.0)				
その他の分類不能の腎炎	176	62	31	13	6	1	1	1	2	2	295	60	355	0.58	2.06
(%)	(59.7)	(21.0)	(10.5)	(4.4)	(2.0)	(0.3)	(0.3)	(0.3)	(0.7)	(0.7)	(100.0)				
多発性嚢胞腎	1,199	464	239	152	91	35	13	9	11	11	2,224	345	2,569	0.71	2.09
(%)	(53.9)	(20.9)	(10.7)	(6.8)	(4.1)	(1.6)	(0.6)	(0.4)	(0.5)	(0.5)	(100.0)				
腎硬化症	2,833	987	580	361	189	65	33	15	16	9	5,088	808	5,896	0.59	1.56
(%)	(55.7)	(19.4)	(11.4)	(7.1)	(3.7)	(1.3)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
悪性高血圧	310	69	49	25	14	4	2	3	1	1	478	77	555	0.50	1.39
(%)	(64.9)	(14.4)	(10.3)	(5.2)	(2.9)	(0.8)	(0.4)	(0.6)	(0.2)	(0.2)	(100.0)				
糖尿病性腎症	11,676	4,334	2,244	1,415	852	248	123	68	64	29	21,053	3,463	24,516	0.58	1.54
(%)	(55.5)	(20.6)	(10.7)	(6.7)	(4.0)	(1.2)	(0.6)	(0.3)	(0.3)	(0.1)	(100.0)				
SLE腎炎	250	115	36	25	16	5	2		2	3	454	83	537	0.63	2.01
(%)	(55.1)	(25.3)	(7.9)	(5.5)	(3.5)	(1.1)	(0.4)		(0.4)	(0.7)	(100.0)				
アミロイド腎	42	23	5	5	3		2				80	18	98	0.54	1.17
(%)	(52.5)	(28.8)	(6.3)	(6.3)	(3.8)		(2.5)				(100.0)				
痛風腎	98	49	17	10	10	5				1	190	37	227	0.79	3.64
(%)	(51.6)	(25.8)	(8.9)	(5.3)	(5.3)	(2.6)				(0.5)	(100.0)				
先天性代謝異常による腎不全	40	8	5	4	1	2					60	13	73	0.43	0.91
(%)	(66.7)	(13.3)	(8.3)	(6.7)	(1.7)	(3.3)					(100.0)				
腎・尿路結核	19	9	4	2	3				1		38	6	44	0.76	1.80
(%)	(50.0)	(23.7)	(10.5)	(5.3)	(7.9)				(2.6)		(100.0)				
腎・尿路結石	56	19	11	5	4	1					96	17	113	0.42	0.72
(%)	(58.3)	(19.8)	(11.5)	(5.2)	(4.2)	(1.0)					(100.0)				
腎・尿路腫瘍	99	35	14	7	5	2		1	2		165	25	190	0.58	1.64
(%)	(60.0)	(21.2)	(8.5)	(4.2)	(3.0)	(1.2)		(0.6)	(1.2)		(100.0)				
閉塞性尿路障害	78	18	19	16	2	2		1			136	15	151	0.50	1.02
(%)	(57.4)	(13.2)	(14.0)	(11.8)	(1.5)	(1.5)		(0.7)			(100.0)				
骨髄腫	18	3	6	2	3	1			1		34	3	37	1.00	2.23
(%)	(52.9)	(8.8)	(17.6)	(5.9)	(8.8)	(2.9)			(2.9)		(100.0)				
腎形成不全	101	24	12	6	3	2	1				149	22	171	0.37	0.89
(%)	(67.8)	(16.1)	(8.1)	(4.0)	(2.0)	(1.3)	(0.7)				(100.0)				
不明	3,208	1,145	544	390	198	65	26	15	14	13	5,618	913	6,531	0.55	1.40
(%)	(57.1)	(20.4)	(9.7)	(6.9)	(3.5)	(1.2)	(0.5)	(0.3)	(0.2)	(0.2)	(100.0)				
再導入	325	115	49	32	17	3	2		2	1	546	87	633	0.46	1.27
(%)	(59.5)	(21.1)	(9.0)	(5.9)	(3.1)	(0.5)	(0.4)		(0.4)	(0.2)	(100.0)				
その他	821	295	149	92	57	22	11	6	4	2	1,459	261	1,720	0.60	1.54
(%)	(56.3)	(20.2)	(10.2)	(6.3)	(3.9)	(1.5)	(0.8)	(0.4)	(0.3)	(0.1)	(100.0)				
合計	33,708	11,651	5,925	3,749	2,104	693	293	162	160	98	58,543	9,831	68,374	0.55	1.50
(%)	(57.6)	(19.9)	(10.1)	(6.4)	(3.6)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
記載なし															
(%)															
総計	33,708	11,651	5,925	3,749	2,104	693	293	162	160	98	58,543	9,831	68,374	0.55	1.50
(%)	(57.6)	(19.9)	(10.1)	(6.4)	(3.6)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D3933##

(C)Japanese Society for Dialysis Therapy

2016年調査