

表 1122 CRP濃度(mg/dL) 医療機関別（腹膜透析患者）

医療機関	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	169	59	29	14	5	3	3	1		1	284	272	556	0.50	1.40
(%)	(59.5)	(20.8)	(10.2)	(4.9)	(1.8)	(1.1)	(1.1)	(0.4)		(0.4)	(100.0)				
私立大学	385	106	71	40	16	9	3	2	5		637	512	1,149	0.55	1.42
(%)	(60.4)	(16.6)	(11.1)	(6.3)	(2.5)	(1.4)	(0.5)	(0.3)	(0.8)		(100.0)				
国立病院機構等	98	29	16	13	10	1	1				168	59	227	0.52	0.96
(%)	(58.3)	(17.3)	(9.5)	(7.7)	(6.0)	(0.6)	(0.6)				(100.0)				
県市町村立・国保	566	192	93	69	45	13	7	5	3	1	994	606	1,600	0.60	1.40
(%)	(56.9)	(19.3)	(9.4)	(6.9)	(4.5)	(1.3)	(0.7)	(0.5)	(0.3)	(0.1)	(100.0)				
JCHO(旧社会保険)等	88	21	21	13	5	2					150	346	496	0.47	0.83
(%)	(58.7)	(14.0)	(14.0)	(8.7)	(3.3)	(1.3)					(100.0)				
厚生連等	178	70	32	30	7	6	2	2	2		329	222	551	0.62	1.38
(%)	(54.1)	(21.3)	(9.7)	(9.1)	(2.1)	(1.8)	(0.6)	(0.6)	(0.6)		(100.0)				
その他公的	476	176	78	59	36	7	1	2	4	3	842	485	1,327	0.59	1.78
(%)	(56.5)	(20.9)	(9.3)	(7.0)	(4.3)	(0.8)	(0.1)	(0.2)	(0.5)	(0.4)	(100.0)				
私立総合	180	71	23	24	16	4	2	3	1	1	325	68	393	0.70	1.80
(%)	(55.4)	(21.8)	(7.1)	(7.4)	(4.9)	(1.2)	(0.6)	(0.9)	(0.3)	(0.3)	(100.0)				
私立	498	170	108	86	39	17	12	2	6	4	942	684	1,626	0.78	1.94
(%)	(52.9)	(18.0)	(11.5)	(9.1)	(4.1)	(1.8)	(1.3)	(0.2)	(0.6)	(0.4)	(100.0)				
私立診療所	280	81	54	31	18	4	3	2	2		475	215	690	0.55	1.31
(%)	(58.9)	(17.1)	(11.4)	(6.5)	(3.8)	(0.8)	(0.6)	(0.4)	(0.4)		(100.0)				
総計	2,918	975	525	379	197	66	34	19	23	10	5,146	3,469	8,615	0.62	1.58
(%)	(56.7)	(18.9)	(10.2)	(7.4)	(3.8)	(1.3)	(0.7)	(0.4)	(0.4)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

##D3944##

(C)Japanese Society for Dialysis Therapy

2016年調査