

表 1128 透析前補正カルシウム濃度(mg/dL) 透析歴(年)別 (透析患者全体)

透析歴	7.0未満	7.0~	7.5~	8.0~	8.4~	8.5~	9.0~	9.5~	10.0~	10.1~	10.5~	11.0~	11.5~	12.0~	合計	記載なし	総計	平均	標準偏差
2年未満	394	574	2,138	6,034	2,609	19,035	17,779	8,493	902	2,065	1,011	361	139	102	61,636	8,902	70,538	8.98	0.72
(%)	(0.6)	(0.9)	(3.5)	(9.8)	(4.2)	(30.9)	(28.8)	(13.8)	(1.5)	(3.4)	(1.6)	(0.6)	(0.2)	(0.2)	(100.0)				
2年~	297	434	1,678	4,928	2,240	19,292	22,491	12,966	1,495	3,334	1,483	518	167	113	71,436	8,732	80,168	9.12	0.70
(%)	(0.4)	(0.6)	(2.3)	(6.9)	(3.1)	(27.0)	(31.5)	(18.2)	(2.1)	(4.7)	(2.1)	(0.7)	(0.2)	(0.2)	(100.0)				
5年~	276	333	1,447	4,019	1,915	16,303	21,944	14,731	1,930	4,642	2,170	712	248	169	70,839	8,306	79,145	9.23	0.73
(%)	(0.4)	(0.5)	(2.0)	(5.7)	(2.7)	(23.0)	(31.0)	(20.8)	(2.7)	(6.6)	(3.1)	(1.0)	(0.4)	(0.2)	(100.0)				
10年~	127	165	717	1,864	852	7,581	10,691	8,226	1,121	2,986	1,510	501	198	86	36,625	3,931	40,556	9.31	0.76
(%)	(0.3)	(0.5)	(2.0)	(5.1)	(2.3)	(20.7)	(29.2)	(22.5)	(3.1)	(8.2)	(4.1)	(1.4)	(0.5)	(0.2)	(100.0)				
15年~	88	116	472	1,136	446	4,044	5,500	4,521	623	1,634	871	281	91	83	19,906	2,188	22,094	9.30	0.79
(%)	(0.4)	(0.6)	(2.4)	(5.7)	(2.2)	(20.3)	(27.6)	(22.7)	(3.1)	(8.2)	(4.4)	(1.4)	(0.5)	(0.4)	(100.0)				
20年~	57	92	285	720	285	2,376	3,102	2,431	370	843	513	136	50	35	11,295	1,189	12,484	9.26	0.80
(%)	(0.5)	(0.8)	(2.5)	(6.4)	(2.5)	(21.0)	(27.5)	(21.5)	(3.3)	(7.5)	(4.5)	(1.2)	(0.4)	(0.3)	(100.0)				
25年~	31	58	189	423	186	1,306	1,711	1,294	184	500	276	72	29	16	6,275	622	6,897	9.24	0.81
(%)	(0.5)	(0.9)	(3.0)	(6.7)	(3.0)	(20.8)	(27.3)	(20.6)	(2.9)	(8.0)	(4.4)	(1.1)	(0.5)	(0.3)	(100.0)				
30年~	10	41	117	240	86	800	928	779	104	275	145	43	21	7	3,596	391	3,987	9.23	0.80
(%)	(0.3)	(1.1)	(3.3)	(6.7)	(2.4)	(22.2)	(25.8)	(21.7)	(2.9)	(7.6)	(4.0)	(1.2)	(0.6)	(0.2)	(100.0)				
35年~	9	16	72	158	48	395	549	404	47	151	74	25	9	7	1,964	188	2,152	9.22	0.81
(%)	(0.5)	(0.8)	(3.7)	(8.0)	(2.4)	(20.1)	(28.0)	(20.6)	(2.4)	(7.7)	(3.8)	(1.3)	(0.5)	(0.4)	(100.0)				
40年~	3	4	22	56	19	151	207	158	19	41	29	18	3	1	731	62	793	9.24	0.79
(%)	(0.4)	(0.5)	(3.0)	(7.7)	(2.6)	(20.7)	(28.3)	(21.6)	(2.6)	(5.6)	(4.0)	(2.5)	(0.4)	(0.1)	(100.0)				
合計	1,292	1,833	7,137	19,578	8,686	71,283	84,902	54,003	6,795	16,471	8,082	2,667	955	619	284,303	34,511	318,814	9.16	0.75
(%)	(0.5)	(0.6)	(2.5)	(6.9)	(3.1)	(25.1)	(29.9)	(19.0)	(2.4)	(5.8)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
不明	2		2	10	4	30	36	34	4	10	14	2	2	4	154	141	295	9.46	1.18
(%)	(1.3)		(1.3)	(6.5)	(2.6)	(19.5)	(23.4)	(22.1)	(2.6)	(6.5)	(9.1)	(1.3)	(1.3)	(2.6)	(100.0)				
記載なし																			
(%)																			
総計	1,294	1,833	7,139	19,588	8,690	71,313	84,938	54,037	6,799	16,481	8,096	2,669	957	623	284,457	34,652	319,109	9.16	0.75
(%)	(0.5)	(0.6)	(2.5)	(6.9)	(3.1)	(25.1)	(29.9)	(19.0)	(2.4)	(5.8)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
平均	6.74	7.27	7.06	6.50	6.25	6.50	7.20	8.32	8.92	9.29	9.83	9.42	9.57	9.27	7.42	6.72	7.35		
標準偏差	7.70	8.69	8.29	7.79	7.37	7.31	7.38	7.72	7.77	7.90	8.17	8.12	8.22	8.13	7.63	7.32	7.60		

数値下のかっこ内は行方向の合計に対する%です。

#D4015#

(C)Japanese Society for Dialysis Therapy

2016年調査