

表 1133 透析前補正カルシウム濃度(mg/dL) 透析前リン濃度(mg/dL)別 (透析患者全体 インタクトPTH値60～240)

透析前リン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
3未満	24	16	93	229	134	1,281	1,663	1,119	145	357	225	84	30	28	5,428	54	5,482	9.29	0.79
(%)	(0.4)	(0.3)	(1.7)	(4.2)	(2.5)	(23.6)	(30.6)	(20.6)	(2.7)	(6.6)	(4.1)	(1.5)	(0.6)	(0.5)	(100.0)				
3.0～	19	28	96	402	187	1,918	2,436	1,400	147	330	179	52	29	12	7,235	49	7,284	9.18	0.68
(%)	(0.3)	(0.4)	(1.3)	(5.6)	(2.6)	(26.5)	(33.7)	(19.4)	(2.0)	(4.6)	(2.5)	(0.7)	(0.4)	(0.2)	(100.0)				
3.5～	41	38	218	719	397	3,594	4,368	2,412	263	636	263	80	27	14	13,070	91	13,161	9.15	0.66
(%)	(0.3)	(0.3)	(1.7)	(5.5)	(3.0)	(27.5)	(33.4)	(18.5)	(2.0)	(4.9)	(2.0)	(0.6)	(0.2)	(0.1)	(100.0)				
4.0～	50	56	329	1,104	588	5,308	6,500	3,597	409	900	371	119	32	15	19,378	99	19,477	9.14	0.65
(%)	(0.3)	(0.3)	(1.7)	(5.7)	(3.0)	(27.4)	(33.5)	(18.6)	(2.1)	(4.6)	(1.9)	(0.6)	(0.2)	(0.1)	(100.0)				
4.5～	55	64	389	1,460	674	6,377	7,595	4,131	463	1,103	450	131	41	23	22,956	138	23,094	9.13	0.65
(%)	(0.2)	(0.3)	(1.7)	(6.4)	(2.9)	(27.8)	(33.1)	(18.0)	(2.0)	(4.8)	(2.0)	(0.6)	(0.2)	(0.1)	(100.0)				
5.0～	46	78	403	1,522	767	6,296	7,339	4,321	492	1,094	441	121	42	15	22,977	133	23,110	9.13	0.65
(%)	(0.2)	(0.3)	(1.8)	(6.6)	(3.3)	(27.4)	(31.9)	(18.8)	(2.1)	(4.8)	(1.9)	(0.5)	(0.2)	(0.1)	(100.0)				
5.5～	55	81	405	1,374	647	5,233	6,021	3,604	436	986	378	106	41	20	19,387	128	19,515	9.13	0.67
(%)	(0.3)	(0.4)	(2.1)	(7.1)	(3.3)	(27.0)	(31.1)	(18.6)	(2.2)	(5.1)	(1.9)	(0.5)	(0.2)	(0.1)	(100.0)				
6.0～	5	24	85	241	133	825	980	615	75	183	72	19	2	3	3,262	17	3,279	9.12	0.69
(%)	(0.2)	(0.7)	(2.6)	(7.4)	(4.1)	(25.3)	(30.0)	(18.9)	(2.3)	(5.6)	(2.2)	(0.6)	(0.1)	(0.1)	(100.0)				
6.1～	32	59	266	811	351	2,951	3,286	1,977	215	600	252	76	22	7	10,905	64	10,969	9.12	0.69
(%)	(0.3)	(0.5)	(2.4)	(7.4)	(3.2)	(27.1)	(30.1)	(18.1)	(2.0)	(5.5)	(2.3)	(0.7)	(0.2)	(0.1)	(100.0)				
6.5～	26	51	260	744	338	2,538	2,659	1,654	215	453	208	62	22	15	9,245	69	9,314	9.10	0.70
(%)	(0.3)	(0.6)	(2.8)	(8.0)	(3.7)	(27.5)	(28.8)	(17.9)	(2.3)	(4.9)	(2.2)	(0.7)	(0.2)	(0.2)	(100.0)				
7.0～	21	48	191	483	168	1,426	1,621	960	119	297	126	37	6	5	5,508	45	5,553	9.09	0.72
(%)	(0.4)	(0.9)	(3.5)	(8.8)	(3.1)	(25.9)	(29.4)	(17.4)	(2.2)	(5.4)	(2.3)	(0.7)	(0.1)	(0.1)	(100.0)				
7.5～	17	33	125	270	115	871	823	574	68	177	75	23	4	6	3,181	41	3,222	9.07	0.75
(%)	(0.5)	(1.0)	(3.9)	(8.5)	(3.6)	(27.4)	(25.9)	(18.0)	(2.1)	(5.6)	(2.4)	(0.7)	(0.1)	(0.2)	(100.0)				
8.0～	12	29	78	173	63	496	504	310	30	110	46	21	3	3	1,878	24	1,902	9.06	0.78
(%)	(0.6)	(1.5)	(4.2)	(9.2)	(3.4)	(26.4)	(26.8)	(16.5)	(1.6)	(5.9)	(2.4)	(1.1)	(0.2)	(0.2)	(100.0)				
8.5～	16	14	55	97	32	283	285	186	23	65	27	9	4	1	1,097	12	1,109	9.02	0.87
(%)	(1.5)	(1.3)	(5.0)	(8.8)	(2.9)	(25.8)	(26.0)	(17.0)	(2.1)	(5.9)	(2.5)	(0.8)	(0.4)	(0.1)	(100.0)				
9.0～	22	36	105	159	57	381	394	277	33	66	32	8	2	1	1,573	22	1,595	8.94	0.85
(%)	(1.4)	(2.3)	(6.7)	(10.1)	(3.6)	(24.2)	(25.0)	(17.6)	(2.1)	(4.2)	(2.0)	(0.5)	(0.1)	(0.1)	(100.0)				
合計	441	655	3,098	9,788	4,651	39,778	46,474	27,137	3,133	7,357	3,145	948	307	168	147,080	986	148,066	9.13	0.68
(%)	(0.3)	(0.4)	(2.1)	(6.7)	(3.2)	(27.0)	(31.6)	(18.5)	(2.1)	(5.0)	(2.1)	(0.6)	(0.2)	(0.1)	(100.0)				
記載なし	2	2	4	4	1	14	27	14	3	10	7	3			91	268	359	9.26	1.05
(%)	(2.2)	(2.2)	(4.4)	(4.4)	(1.1)	(15.4)	(29.7)	(15.4)	(3.3)	(11.0)	(7.7)	(3.3)			(100.0)				
総計	443	657	3,102	9,792	4,652	39,792	46,501	27,151	3,136	7,367	3,152	951	307	168	147,171	1,254	148,425	9.13	0.68
(%)	(0.3)	(0.4)	(2.1)	(6.7)	(3.2)	(27.0)	(31.6)	(18.4)	(2.1)	(5.0)	(2.1)	(0.6)	(0.2)	(0.1)	(100.0)				
平均	5.52	5.88	5.61	5.38	5.26	5.15	5.08	5.12	5.15	5.17	5.09	5.03	4.85	4.70	5.16	5.30	5.16		
標準偏差	1.84	1.81	1.61	1.41	1.36	1.33	1.30	1.35	1.37	1.40	1.48	1.56	1.47	1.82	1.36	1.57	1.36		

数値下のかっこ内は行方向の合計に対する%です。
##D4022##
(C)Japanese Society for Dialysis Therapy
2016年調査