

表 1134 透析前補正カルシウム濃度(mg/dL) 性別 (施設血液透析患者 透析回数週3回)

性別	7.0未満	7.0~	7.5~	8.0~	8.4~	8.5~	9.0~	9.5~	10.0~	10.1~	10.5~	11.0~	11.5~	12.0~	合計	記載なし	総計	平均	標準偏差
男性	607	908	3,464	9,889	4,354	34,652	38,225	22,800	2,788	6,741	3,259	1,051	361	249	129,348	6,102	135,450	9.11	0.74
(%)	(0.5)	(0.7)	(2.7)	(7.6)	(3.4)	(26.8)	(29.6)	(17.6)	(2.2)	(5.2)	(2.5)	(0.8)	(0.3)	(0.2)	(100.0)				
女性	348	395	1,460	3,811	1,807	15,849	21,660	14,845	1,954	4,784	2,503	870	342	223	70,851	3,234	74,085	9.25	0.76
(%)	(0.5)	(0.6)	(2.1)	(5.4)	(2.6)	(22.4)	(30.6)	(21.0)	(2.8)	(6.8)	(3.5)	(1.2)	(0.5)	(0.3)	(100.0)				
合計	955	1,303	4,924	13,700	6,161	50,501	59,885	37,645	4,742	11,525	5,762	1,921	703	472	200,199	9,336	209,535	9.16	0.75
(%)	(0.5)	(0.7)	(2.5)	(6.8)	(3.1)	(25.2)	(29.9)	(18.8)	(2.4)	(5.8)	(2.9)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし																1	1		
総計	955	1,303	4,924	13,700	6,161	50,501	59,885	37,645	4,742	11,525	5,762	1,921	703	472	200,199	9,337	209,536	9.16	0.75
(%)	(0.5)	(0.7)	(2.5)	(6.8)	(3.1)	(25.2)	(29.9)	(18.8)	(2.4)	(5.8)	(2.9)	(1.0)	(0.4)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

#D4030##

(C)Japanese Society for Dialysis Therapy

2016年調査