

表 1137 透析前補正カルシウム濃度(mg/dL) 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	254	347	1,281	3,371	1,415	12,473	16,073	11,308	1,536	3,735	1,862	614	228	153	54,650	2,522	57,172	9.23	0.76
(%)	(0.5)	(0.6)	(2.3)	(6.2)	(2.6)	(22.8)	(29.4)	(20.7)	(2.8)	(6.8)	(3.4)	(1.1)	(0.4)	(0.3)	(100.0)				
慢性腎盂腎炎	5	15	43	93	51	391	529	397	51	131	74	17	4	3	1,804	64	1,868	9.26	0.75
(%)	(0.3)	(0.8)	(2.4)	(5.2)	(2.8)	(21.7)	(29.3)	(22.0)	(2.8)	(7.3)	(4.1)	(0.9)	(0.2)	(0.2)	(100.0)				
急速進行性糸球体腎炎	10	8	33	106	48	396	538	285	36	82	40	16	4	3	1,605	82	1,687	9.15	0.72
(%)	(0.6)	(0.5)	(2.1)	(6.6)	(3.0)	(24.7)	(33.5)	(17.8)	(2.2)	(5.1)	(2.5)	(1.0)	(0.2)	(0.2)	(100.0)				
妊娠腎/妊娠中毒症	6	3	21	57	18	143	228	185	25	78	29	9	5	1	808	41	849	9.28	0.80
(%)	(0.7)	(0.4)	(2.6)	(7.1)	(2.2)	(17.7)	(28.2)	(22.9)	(3.1)	(9.7)	(3.6)	(1.1)	(0.6)	(0.1)	(100.0)				
その他の分類不能の腎炎	7	15	20	44	33	199	266	169	23	51	27	8	1	1	864	38	902	9.15	0.78
(%)	(0.8)	(1.7)	(2.3)	(5.1)	(3.8)	(23.0)	(30.8)	(19.6)	(2.7)	(5.9)	(3.1)	(0.9)	(0.1)	(0.1)	(100.0)				
多発性嚢胞腎	33	37	140	392	173	1,678	2,215	1,545	176	509	243	67	32	15	7,255	293	7,548	9.24	0.74
(%)	(0.5)	(0.5)	(1.9)	(5.4)	(2.4)	(23.1)	(30.5)	(21.3)	(2.4)	(7.0)	(3.3)	(0.9)	(0.4)	(0.2)	(100.0)				
腎硬化症	85	121	471	1,374	665	5,358	6,389	3,816	445	1,064	537	168	60	52	20,605	913	21,518	9.15	0.72
(%)	(0.4)	(0.6)	(2.3)	(6.7)	(3.2)	(26.0)	(31.0)	(18.5)	(2.2)	(5.2)	(2.6)	(0.8)	(0.3)	(0.3)	(100.0)				
悪性高血圧	12	5	59	115	48	423	484	336	44	85	52	8	6	2	1,679	98	1,777	9.14	0.75
(%)	(0.7)	(0.3)	(3.5)	(6.8)	(2.9)	(25.2)	(28.8)	(20.0)	(2.6)	(5.1)	(3.1)	(0.5)	(0.4)	(0.1)	(100.0)				
糖尿病性腎症	405	572	2,133	6,099	2,829	21,944	24,233	13,805	1,699	3,966	1,961	706	253	165	80,770	3,698	84,468	9.11	0.74
(%)	(0.5)	(0.7)	(2.6)	(7.6)	(3.5)	(27.2)	(30.0)	(17.1)	(2.1)	(4.9)	(2.4)	(0.9)	(0.3)	(0.2)	(100.0)				
SLE腎炎	5	6	31	77	33	287	393	273	31	107	45	12	4	1	1,305	65	1,370	9.23	0.74
(%)	(0.4)	(0.5)	(2.4)	(5.9)	(2.5)	(22.0)	(30.1)	(20.9)	(2.4)	(8.2)	(3.4)	(0.9)	(0.3)	(0.1)	(100.0)				
アミロイド腎	2		5	17	7	69	72	63	4	17	7	3		1	267	13	280	9.20	0.70
(%)	(0.7)		(1.9)	(6.4)	(2.6)	(25.8)	(27.0)	(23.6)	(1.5)	(6.4)	(2.6)	(1.1)		(0.4)	(100.0)				
痛風腎	2	5	11	65	13	154	196	129	15	48	20	4	3	2	667	33	700	9.19	0.76
(%)	(0.3)	(0.7)	(1.6)	(9.7)	(1.9)	(23.1)	(29.4)	(19.3)	(2.2)	(7.2)	(3.0)	(0.6)	(0.4)	(0.3)	(100.0)				
先天性代謝異常による腎不全	2	4	6	17	3	29	52	30	4	7	5	1	2		162	7	169	9.10	0.82
(%)	(1.2)	(2.5)	(3.7)	(10.5)	(1.9)	(17.9)	(32.1)	(18.5)	(2.5)	(4.3)	(3.1)	(0.6)	(1.2)		(100.0)				
腎・尿路結核		2	1	7	1	21	32	17	2	10	5			2	100	6	106	9.29	0.81
(%)		(2.0)	(1.0)	(7.0)	(1.0)	(21.0)	(32.0)	(17.0)	(2.0)	(10.0)	(5.0)			(2.0)	(100.0)				
腎・尿路結石	1	3	9	20	10	99	110	62	7	20	21	3	1	1	367	17	384	9.20	0.77
(%)	(0.3)	(0.8)	(2.5)	(5.4)	(2.7)	(27.0)	(30.0)	(16.9)	(1.9)	(5.4)	(5.7)	(0.8)	(0.3)	(0.3)	(100.0)				
腎・尿路腫瘍	1	7	14	47	15	179	179	109	8	42	16	4	2	3	626	18	644	9.14	0.74
(%)	(0.2)	(1.1)	(2.2)	(7.5)	(2.4)	(28.6)	(28.6)	(17.4)	(1.3)	(6.7)	(2.6)	(0.6)	(0.3)	(0.5)	(100.0)				
閉塞性尿路障害	2	5	13	23	12	112	132	77	15	29	13	6	1	1	441	24	465	9.19	0.74
(%)	(0.5)	(1.1)	(2.9)	(5.2)	(2.7)	(25.4)	(29.9)	(17.5)	(3.4)	(6.6)	(2.9)	(1.4)	(0.2)	(0.2)	(100.0)				
骨髄腫	1	2	11	16	4	59	58	26	4	13	4	1	1		200	13	213	9.05	0.75
(%)	(0.5)	(1.0)	(5.5)	(8.0)	(2.0)	(29.5)	(29.0)	(13.0)	(2.0)	(6.5)	(2.0)	(0.5)	(0.5)		(100.0)				
腎形成不全	3	2	11	23	13	81	103	69	10	23	10	5	1	1	355	13	368	9.18	0.83
(%)	(0.8)	(0.6)	(3.1)	(6.5)	(3.7)	(22.8)	(29.0)	(19.4)	(2.8)	(6.5)	(2.8)	(1.4)	(0.3)	(0.3)	(100.0)				
不明	74	99	453	1,321	580	4,877	5,769	3,708	440	1,137	580	206	76	53	19,373	1,105	20,478	9.18	0.75
(%)	(0.4)	(0.5)	(2.3)	(6.8)	(3.0)	(25.2)	(29.8)	(19.1)	(2.3)	(5.9)	(3.0)	(1.1)	(0.4)	(0.3)	(100.0)				
再導入	10	14	26	75	31	259	301	239	32	74	41	10	1	2	1,115	49	1,164	9.18	0.78
(%)	(0.9)	(1.3)	(2.3)	(6.7)	(2.8)	(23.2)	(27.0)	(21.4)	(2.9)	(6.6)	(3.7)	(0.9)	(0.1)	(0.2)	(100.0)				
その他	35	31	132	341	159	1,270	1,533	997	135	297	170	53	18	10	5,181	225	5,406	9.17	0.78
(%)	(0.7)	(0.6)	(2.5)	(6.6)	(3.1)	(24.5)	(29.6)	(19.2)	(2.6)	(5.7)	(3.3)	(1.0)	(0.3)	(0.2)	(100.0)				
合計	955	1,303	4,924	13,700	6,161	50,501	59,885	37,645	4,742	11,525	5,762	1,921	703	472	200,199	9,337	209,536	9.16	0.75
(%)	(0.5)	(0.7)	(2.5)	(6.8)	(3.1)	(25.2)	(29.9)	(18.8)	(2.4)	(5.8)	(2.9)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし																			
(%)																			
総計	955	1,303	4,924	13,700	6,161	50,501	59,885	37,645	4,742	11,525	5,762	1,921	703	472	200,199	9,337	209,536	9.16	0.75
(%)	(0.5)	(0.7)	(2.5)	(6.8)	(3.1)	(25.2)	(29.9)	(18.8)	(2.4)	(5.8)	(2.9)	(1.0)	(0.4)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4033##

(C)Japanese Society for Dialysis Therapy

2016年調査