

表 1141 透析前補正カルシウム濃度(mg/dL) 透析歴(年)別 (血液透析濾過患者 透析回数週3回)

透析歴	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
2年未満	34	87	351	988	416	3,122	2,781	1,291	135	297	147	48	14	18	9,729	289	10,018	8.96	0.69
(%)	(0.3)	(0.9)	(3.6)	(10.2)	(4.3)	(32.1)	(28.6)	(13.3)	(1.4)	(3.1)	(1.5)	(0.5)	(0.1)	(0.2)	(100.0)				
2年～	52	95	380	1,146	510	4,326	4,888	2,737	312	633	263	83	34	14	15,473	465	15,938	9.09	0.67
(%)	(0.3)	(0.6)	(2.5)	(7.4)	(3.3)	(28.0)	(31.6)	(17.7)	(2.0)	(4.1)	(1.7)	(0.5)	(0.2)	(0.1)	(100.0)				
5年～	56	78	378	1,107	496	4,013	5,288	3,414	433	1,043	472	144	39	34	16,995	587	17,582	9.19	0.72
(%)	(0.3)	(0.5)	(2.2)	(6.5)	(2.9)	(23.6)	(31.1)	(20.1)	(2.5)	(6.1)	(2.8)	(0.8)	(0.2)	(0.2)	(100.0)				
10年～	27	38	216	525	235	2,153	2,897	2,129	295	733	360	117	45	15	9,785	331	10,116	9.27	0.74
(%)	(0.3)	(0.4)	(2.2)	(5.4)	(2.4)	(22.0)	(29.6)	(21.8)	(3.0)	(7.5)	(3.7)	(1.2)	(0.5)	(0.2)	(100.0)				
15年～	21	37	155	370	137	1,289	1,650	1,288	185	447	197	65	19	13	5,873	185	6,058	9.24	0.76
(%)	(0.4)	(0.6)	(2.6)	(6.3)	(2.3)	(21.9)	(28.1)	(21.9)	(3.2)	(7.6)	(3.4)	(1.1)	(0.3)	(0.2)	(100.0)				
20年～	16	25	106	281	91	803	1,036	814	120	255	144	39	11	5	3,746	111	3,857	9.22	0.76
(%)	(0.4)	(0.7)	(2.8)	(7.5)	(2.4)	(21.4)	(27.7)	(21.7)	(3.2)	(6.8)	(3.8)	(1.0)	(0.3)	(0.1)	(100.0)				
25年～	12	15	71	159	73	483	649	471	53	166	85	21	6	6	2,270	75	2,345	9.21	0.78
(%)	(0.5)	(0.7)	(3.1)	(7.0)	(3.2)	(21.3)	(28.6)	(20.7)	(2.3)	(7.3)	(3.7)	(0.9)	(0.3)	(0.3)	(100.0)				
30年～	4	12	37	84	34	320	341	280	32	104	53	17	3	3	1,324	43	1,367	9.22	0.77
(%)	(0.3)	(0.9)	(2.8)	(6.3)	(2.6)	(24.2)	(25.8)	(21.1)	(2.4)	(7.9)	(4.0)	(1.3)	(0.2)	(0.2)	(100.0)				
35年～	3	8	22	64	19	169	216	154	15	54	26	6	4	1	761	19	780	9.18	0.78
(%)	(0.4)	(1.1)	(2.9)	(8.4)	(2.5)	(22.2)	(28.4)	(20.2)	(2.0)	(7.1)	(3.4)	(0.8)	(0.5)	(0.1)	(100.0)				
40年～		1	9	26	7	62	80	56	4	18	10	6		1	280	7	287	9.20	0.76
(%)		(0.4)	(3.2)	(9.3)	(2.5)	(22.1)	(28.6)	(20.0)	(1.4)	(6.4)	(3.6)	(2.1)		(0.4)	(100.0)				
合計	225	396	1,725	4,750	2,018	16,740	19,826	12,634	1,584	3,750	1,757	546	175	110	66,236	2,112	68,348	9.15	0.72
(%)	(0.3)	(0.6)	(2.6)	(7.2)	(3.0)	(25.3)	(29.9)	(19.1)	(2.4)	(5.7)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
不明				1	1	5	2	3		1					13	13	26	9.02	0.54
(%)				(7.7)	(7.7)	(38.5)	(15.4)	(23.1)		(7.7)					(100.0)				
記載なし																			
(%)																			
総計	225	396	1,725	4,751	2,019	16,745	19,828	12,637	1,584	3,751	1,757	546	175	110	66,249	2,125	68,374	9.15	0.72
(%)	(0.3)	(0.6)	(2.6)	(7.2)	(3.0)	(25.3)	(29.9)	(19.1)	(2.4)	(5.7)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
平均	9.39	9.17	9.07	8.49	7.93	8.22	8.87	10.00	10.39	11.07	11.58	11.32	10.81	10.30	9.14	9.10	9.14		
標準偏差	8.68	9.59	9.07	8.79	8.31	8.30	8.25	8.50	8.11	8.65	8.90	8.77	8.28	9.08	8.48	8.24	8.47		

数値下のかっこ内は行方向の合計に対する%です。
##D4042##
(C)Japanese Society for Dialysis Therapy
2016年調査