

表 1152 透析前リン濃度(mg/dL) 性別 (透析患者全体)

性別	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
男性	7,433	9,007	15,957	22,856	27,605	27,851	24,038	4,138	14,057	12,506	7,740	4,816	2,920	1,853	2,920	185,697	20,226	205,923	5.23	1.45
(%)	(4.0)	(4.9)	(8.6)	(12.3)	(14.9)	(15.0)	(12.9)	(2.2)	(7.6)	(6.7)	(4.2)	(2.6)	(1.6)	(1.0)	(1.6)	(100.0)				
女性	4,771	5,225	9,022	13,081	15,526	15,568	13,221	2,180	7,537	6,355	3,887	2,219	1,314	807	1,289	102,002	11,183	113,185	5.14	1.41
(%)	(4.7)	(5.1)	(8.8)	(12.8)	(15.2)	(15.3)	(13.0)	(2.1)	(7.4)	(6.2)	(3.8)	(2.2)	(1.3)	(0.8)	(1.3)	(100.0)				
合計	12,204	14,232	24,979	35,937	43,131	43,419	37,259	6,318	21,594	18,861	11,627	7,035	4,234	2,660	4,209	287,699	31,409	319,108	5.20	1.44
(%)	(4.2)	(4.9)	(8.7)	(12.5)	(15.0)	(15.1)	(13.0)	(2.2)	(7.5)	(6.6)	(4.0)	(2.4)	(1.5)	(0.9)	(1.5)	(100.0)				
記載なし																	1	1		
(%)																				
総計	12,204	14,232	24,979	35,937	43,131	43,419	37,259	6,318	21,594	18,861	11,627	7,035	4,234	2,660	4,209	287,699	31,410	319,109	5.20	1.44
(%)	(4.2)	(4.9)	(8.7)	(12.5)	(15.0)	(15.1)	(13.0)	(2.2)	(7.5)	(6.6)	(4.0)	(2.4)	(1.5)	(0.9)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
##D4113##
(C)Japanese Society for Dialysis Therapy
2016年調査