

表 1155 透析前リン濃度(mg/dL) 原疾患別 (透析患者全体)

原疾患	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	3,080	3,685	6,741	10,141	12,578	13,087	11,359	1,929	6,491	5,549	3,323	2,017	1,166	732	1,179	83,057	8,726	91,783	5.24	1.40
(%)	(3.7)	(4.4)	(8.1)	(12.2)	(15.1)	(15.8)	(13.7)	(2.3)	(7.8)	(6.7)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
慢性腎盂腎炎	124	142	222	349	403	377	335	61	181	172	109	70	31	33	41	2,650	246	2,896	5.18	1.47
(%)	(4.7)	(5.4)	(8.4)	(13.2)	(15.2)	(14.2)	(12.6)	(2.3)	(6.8)	(6.5)	(4.1)	(2.6)	(1.2)	(1.2)	(1.5)	(100.0)				
急速進行性糸球体腎炎	142	142	229	311	357	313	261	39	150	124	71	46	23	21	39	2,268	270	2,538	5.02	1.49
(%)	(6.3)	(6.3)	(10.1)	(13.7)	(15.7)	(13.8)	(11.5)	(1.7)	(6.6)	(5.5)	(3.1)	(2.0)	(1.0)	(0.9)	(1.7)	(100.0)				
妊娠腎/妊娠中毒症	45	45	100	166	206	181	206	25	119	107	63	46	26	10	27	1,372	111	1,483	5.38	1.44
(%)	(3.3)	(3.3)	(7.3)	(12.1)	(15.0)	(13.2)	(15.0)	(1.8)	(8.7)	(7.8)	(4.6)	(3.4)	(1.9)	(0.7)	(2.0)	(100.0)				
その他の分類不能の腎炎	44	70	135	142	186	180	177	25	106	91	56	43	31	17	25	1,328	126	1,454	5.32	1.51
(%)	(3.3)	(5.3)	(10.2)	(10.7)	(14.0)	(13.6)	(13.3)	(1.9)	(8.0)	(6.9)	(4.2)	(3.2)	(2.3)	(1.3)	(1.9)	(100.0)				
多発性嚢胞腎	304	359	739	1,218	1,645	1,709	1,521	214	903	767	485	257	131	90	133	10,475	996	11,471	5.32	1.34
(%)	(2.9)	(3.4)	(7.1)	(11.6)	(15.7)	(16.3)	(14.5)	(2.0)	(8.6)	(7.3)	(4.6)	(2.5)	(1.3)	(0.9)	(1.3)	(100.0)				
腎硬化症	1,392	1,548	2,778	3,833	4,443	4,228	3,545	557	2,005	1,725	1,038	611	364	226	314	28,607	3,043	31,650	5.08	1.41
(%)	(4.9)	(5.4)	(9.7)	(13.4)	(15.5)	(14.8)	(12.4)	(1.9)	(7.0)	(6.0)	(3.6)	(2.1)	(1.3)	(0.8)	(1.1)	(100.0)				
悪性高血圧	92	112	212	293	332	354	291	50	190	188	101	79	46	30	54	2,424	297	2,721	5.33	1.54
(%)	(3.8)	(4.6)	(8.7)	(12.1)	(13.7)	(14.6)	(12.0)	(2.1)	(7.8)	(7.8)	(4.2)	(3.3)	(1.9)	(1.2)	(2.2)	(100.0)				
糖尿病性腎症	4,817	5,698	9,899	13,955	16,492	16,572	14,094	2,465	8,231	7,349	4,591	2,835	1,784	1,075	1,689	111,546	12,111	123,657	5.20	1.46
(%)	(4.3)	(5.1)	(8.9)	(12.5)	(14.8)	(14.9)	(12.6)	(2.2)	(7.4)	(6.6)	(4.1)	(2.5)	(1.6)	(1.0)	(1.5)	(100.0)				
SLE腎炎	75	95	158	264	292	299	278	50	153	115	80	53	24	22	27	1,985	237	2,222	5.23	1.42
(%)	(3.8)	(4.8)	(8.0)	(13.3)	(14.7)	(15.1)	(14.0)	(2.5)	(7.7)	(5.8)	(4.0)	(2.7)	(1.2)	(1.1)	(1.4)	(100.0)				
アミロイド腎	14	23	46	49	52	60	51	6	29	23	23	4	5	5	1	391	47	438	5.09	1.34
(%)	(3.6)	(5.9)	(11.8)	(12.5)	(13.3)	(15.3)	(13.0)	(1.5)	(7.4)	(5.9)	(5.9)	(1.0)	(1.3)	(1.3)	(0.3)	(100.0)				
痛風腎	38	46	64	99	146	141	122	35	73	74	52	22	14	6	24	956	100	1,056	5.35	1.53
(%)	(4.0)	(4.8)	(6.7)	(10.4)	(15.3)	(14.7)	(12.8)	(3.7)	(7.6)	(7.7)	(5.4)	(2.3)	(1.5)	(0.6)	(2.5)	(100.0)				
先天性代謝異常による腎不全	7	6	21	31	26	50	26	3	17	22	13	11	10	7	6	256	25	281	5.59	1.61
(%)	(2.7)	(2.3)	(8.2)	(12.1)	(10.2)	(19.5)	(10.2)	(1.2)	(6.6)	(8.6)	(5.1)	(4.3)	(3.9)	(2.7)	(2.3)	(100.0)				
腎・尿路結核	5	6	11	26	25	19	19	5	9	13	4	2	5		1	150	18	168	5.16	1.46
(%)	(3.3)	(4.0)	(7.3)	(17.3)	(16.7)	(12.7)	(12.7)	(3.3)	(6.0)	(8.7)	(2.7)	(1.3)	(3.3)		(0.7)	(100.0)				
腎・尿路結石	19	26	43	69	83	90	53	9	46	27	22	12	8	5	6	518	60	578	5.16	1.37
(%)	(3.7)	(5.0)	(8.3)	(13.3)	(16.0)	(17.4)	(10.2)	(1.7)	(8.9)	(5.2)	(4.2)	(2.3)	(1.5)	(1.0)	(1.2)	(100.0)				
腎・尿路腫瘍	33	45	71	116	117	135	107	24	76	46	39	19	17	6	8	859	100	959	5.20	1.35
(%)	(3.8)	(5.2)	(8.3)	(13.5)	(13.6)	(15.7)	(12.5)	(2.8)	(8.8)	(5.4)	(4.5)	(2.2)	(2.0)	(0.7)	(0.9)	(100.0)				
閉塞性尿路障害	26	42	73	96	94	91	75	9	44	34	27	21	9	5	14	660	83	743	5.13	1.51
(%)	(3.9)	(6.4)	(11.1)	(14.5)	(14.2)	(13.8)	(11.4)	(1.4)	(6.7)	(5.2)	(4.1)	(3.2)	(1.4)	(0.8)	(2.1)	(100.0)				
骨髄腫	18	26	41	40	38	34	32	4	21	6	6	6	5	2	4	283	40	323	4.82	1.45
(%)	(6.4)	(9.2)	(14.5)	(14.1)	(13.4)	(12.0)	(11.3)	(1.4)	(7.4)	(2.1)	(2.1)	(2.1)	(1.8)	(0.7)	(1.4)	(100.0)				
腎形成不全	12	24	33	65	69	94	82	14	46	63	38	13	14	14	20	601	63	664	5.64	1.61
(%)	(2.0)	(4.0)	(5.5)	(10.8)	(11.5)	(15.6)	(13.6)	(2.3)	(7.7)	(10.5)	(6.3)	(2.2)	(2.3)	(2.3)	(3.3)	(100.0)				
不明	1,476	1,576	2,545	3,443	4,200	4,036	3,488	609	2,024	1,744	1,072	595	383	261	412	27,864	3,548	31,412	5.13	1.46
(%)	(5.3)	(5.7)	(9.1)	(12.4)	(15.1)	(14.5)	(12.5)	(2.2)	(7.3)	(6.3)	(3.8)	(2.1)	(1.4)	(0.9)	(1.5)	(100.0)				
再導入	53	76	129	221	271	286	263	37	149	157	106	75	41	25	43	1,932	242	2,174	5.47	1.50
(%)	(2.7)	(3.9)	(6.7)	(11.4)	(14.0)	(14.8)	(13.6)	(1.9)	(7.7)	(8.1)	(5.5)	(3.9)	(2.1)	(1.3)	(2.2)	(100.0)				
その他	388	440	689	1,010	1,076	1,083	874	148	531	465	308	198	97	68	142	7,517	921	8,438	5.14	1.50
(%)	(5.2)	(5.9)	(9.2)	(13.4)	(14.3)	(14.4)	(11.6)	(2.0)	(7.1)	(6.2)	(4.1)	(2.6)	(1.3)	(0.9)	(1.9)	(100.0)				
合計	12,204	14,232	24,979	35,937	43,131	43,419	37,259	6,318	21,594	18,861	11,627	7,035	4,234	2,660	4,209	287,699	31,410	319,109	5.20	1.44
(%)	(4.2)	(4.9)	(8.7)	(12.5)	(15.0)	(15.1)	(13.0)	(2.2)	(7.5)	(6.6)	(4.0)	(2.4)	(1.5)	(0.9)	(1.5)	(100.0)				
記載なし																				
(%)																				
総計	12,204	14,232	24,979	35,937	43,131	43,419	37,259	6,318	21,594	18,861	11,627	7,035	4,234	2,660	4,209	287,699	31,410	319,109	5.20	1.44
(%)	(4.2)	(4.9)	(8.7)	(12.5)	(15.0)	(15.1)	(13.0)	(2.2)	(7.5)	(6.6)	(4.0)	(2.4)	(1.5)	(0.9)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
##D4116##
(C)Japanese Society for Dialysis Therapy
2016年調査