

表 1160 透析前リソルビン濃度(mg/dL) 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	3未満	3.0~	3.5~	4.0~	4.5~	5.0~	5.5~	6.0~	6.1~	6.5~	7.0~	7.5~	8.0~	8.5~	9.0~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,245	2,567	4,674	6,861	8,408	8,710	7,378	1,279	4,258	3,624	2,167	1,312	746	459	780	55,468	1,704	57,172	5.20	1.41
(%)	(4.0)	(4.6)	(8.4)	(12.4)	(15.2)	(15.7)	(13.3)	(2.3)	(7.7)	(6.5)	(3.9)	(2.4)	(1.3)	(0.8)	(1.4)	(100.0)				
慢性腎孟腎炎	87	99	153	235	289	263	231	36	132	116	69	46	23	21	28	1,828	40	1,868	5.17	1.47
(%)	(4.8)	(5.4)	(8.4)	(12.9)	(15.8)	(14.4)	(12.6)	(2.0)	(7.2)	(6.3)	(3.8)	(2.5)	(1.3)	(1.1)	(1.5)	(100.0)				
急速進行性糸球体腎炎	112	99	164	228	256	216	191	27	101	81	47	34	18	15	31	1,620	67	1,687	5.01	1.53
(%)	(6.9)	(6.1)	(10.1)	(14.1)	(15.8)	(13.3)	(11.8)	(1.7)	(6.2)	(5.0)	(2.9)	(2.1)	(1.1)	(0.9)	(1.9)	(100.0)				
妊娠腎/妊娠中毒症	24	23	63	102	120	111	132	13	70	57	40	32	19	5	15	826	23	849	5.41	1.44
(%)	(2.9)	(2.8)	(7.6)	(12.3)	(14.5)	(13.4)	(16.0)	(1.6)	(8.5)	(6.9)	(4.8)	(3.9)	(2.3)	(0.6)	(1.8)	(100.0)				
その他の分類不能の腎炎	27	51	92	97	130	128	110	15	52	57	37	30	23	11	14	874	28	902	5.27	1.50
(%)	(3.1)	(5.8)	(10.5)	(11.1)	(14.9)	(14.6)	(12.6)	(1.7)	(5.9)	(6.5)	(4.2)	(3.4)	(2.6)	(1.3)	(1.6)	(100.0)				
多発性囊胞腎	224	266	531	865	1,195	1,197	1,036	148	624	521	309	193	81	63	84	7,337	211	7,548	5.28	1.34
(%)	(3.1)	(3.6)	(7.2)	(11.8)	(16.3)	(16.3)	(14.1)	(2.0)	(8.5)	(7.1)	(4.2)	(2.6)	(1.1)	(0.9)	(1.1)	(100.0)				
腎硬化症	1,064	1,124	2,020	2,775	3,278	3,069	2,603	401	1,470	1,244	753	449	261	160	213	20,884	634	21,518	5.07	1.40
(%)	(5.1)	(5.4)	(9.7)	(13.3)	(15.7)	(14.7)	(12.5)	(1.9)	(7.0)	(6.0)	(3.6)	(2.1)	(1.2)	(0.8)	(1.0)	(100.0)				
悪性高血圧	67	82	163	209	240	247	198	39	112	139	68	51	33	22	35	1,705	72	1,777	5.28	1.54
(%)	(3.9)	(4.8)	(9.6)	(12.3)	(14.1)	(14.5)	(11.6)	(2.3)	(6.6)	(8.2)	(4.0)	(3.0)	(1.9)	(1.3)	(2.1)	(100.0)				
糖尿病性腎症	3,696	4,268	7,412	10,242	11,987	12,211	10,211	1,786	6,025	5,284	3,298	2,025	1,248	775	1,212	81,680	2,788	84,468	5.18	1.46
(%)	(4.5)	(5.2)	(9.1)	(12.5)	(14.7)	(14.9)	(12.5)	(2.2)	(7.4)	(6.5)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
SLE腎炎	46	60	109	172	196	195	185	30	104	82	52	37	15	13	21	1,317	53	1,370	5.25	1.42
(%)	(3.5)	(4.6)	(8.3)	(13.1)	(14.9)	(14.8)	(14.0)	(2.3)	(7.9)	(6.2)	(3.9)	(2.8)	(1.1)	(1.0)	(1.6)	(100.0)				
アミロイド腎	12	19	34	27	36	41	37	4	20	16	14	4	2	5	271	9	280	5.05	1.37	
(%)	(4.4)	(7.0)	(12.5)	(10.0)	(13.3)	(15.1)	(13.7)	(1.5)	(7.4)	(5.9)	(5.2)	(1.5)	(0.7)	(1.8)	(100.0)					
痛風腎	28	34	53	71	99	105	80	25	49	51	33	16	13	1	17	675	25	700	5.31	1.54
(%)	(4.1)	(5.0)	(7.9)	(10.5)	(14.7)	(15.6)	(11.9)	(3.7)	(7.3)	(7.6)	(4.9)	(2.4)	(1.9)	(0.1)	(2.5)	(100.0)				
先天性代謝異常による腎不全	6	5	16	16	17	31	17	2	12	12	9	3	5	6	6	163	6	169	5.56	1.73
(%)	(3.7)	(3.1)	(9.8)	(9.8)	(10.4)	(19.0)	(10.4)	(1.2)	(7.4)	(7.4)	(5.5)	(1.8)	(3.1)	(3.7)	(3.7)	(100.0)				
腎・尿路結核	3	5	5	20	15	11	16	3	6	10	3	1	1	1	1	100	6	106	5.15	1.47
(%)	(3.0)	(5.0)	(5.0)	(20.0)	(15.0)	(11.0)	(16.0)	(3.0)	(6.0)	(10.0)	(3.0)	(1.0)	(1.0)	(1.0)	(1.0)	(100.0)				
腎・尿路結石	14	20	28	47	59	64	40	8	35	19	18	7	6	3	3	371	13	384	5.17	1.36
(%)	(3.8)	(5.4)	(7.5)	(12.7)	(15.9)	(17.3)	(10.8)	(2.2)	(9.4)	(5.1)	(4.9)	(1.9)	(1.6)	(0.8)	(0.8)	(100.0)				
腎・尿路腫瘍	23	33	54	82	90	93	83	17	54	32	26	14	14	5	7	627	17	644	5.21	1.36
(%)	(3.7)	(5.3)	(8.6)	(13.1)	(14.4)	(14.8)	(13.2)	(2.7)	(8.6)	(5.1)	(4.1)	(2.2)	(2.2)	(0.8)	(1.1)	(100.0)				
閉塞性尿路障害	17	27	48	62	68	60	54	7	27	22	20	14	5	4	9	444	21	465	5.14	1.49
(%)	(3.8)	(6.1)	(10.8)	(14.0)	(15.3)	(13.5)	(12.2)	(1.6)	(6.1)	(5.0)	(4.5)	(3.2)	(1.1)	(0.9)	(2.0)	(100.0)				
骨髄腫	14	15	31	29	24	27	24	3	12	5	2	5	3	2	3	199	14	213	4.80	1.47
(%)	(7.0)	(7.5)	(15.6)	(14.6)	(12.1)	(13.6)	(12.1)	(1.5)	(6.0)	(2.5)	(1.0)	(2.5)	(1.5)	(1.0)	(1.5)	(100.0)				
腎形成不全	8	11	20	41	43	66	46	5	29	44	19	6	6	8	10	362	6	368	5.58	1.58
(%)	(2.2)	(3.0)	(5.5)	(11.3)	(11.9)	(18.2)	(12.7)	(1.4)	(8.0)	(12.2)	(5.2)	(1.7)	(1.7)	(2.2)	(2.8)	(100.0)				
不明	1,112	1,148	1,801	2,424	2,949	2,858	2,422	418	1,398	1,218	736	409	252	169	281	19,955	883	20,478	5.10	1.46
(%)	(5.7)	(5.9)	(9.2)	(12.4)	(15.0)	(14.6)	(12.4)	(2.1)	(7.1)	(6.2)	(3.8)	(2.1)	(1.3)	(0.9)	(1.4)	(100.0)				
再導入	32	49	80	131	152	169	142	19	85	94	56	44	26	17	29	1,125	39	1,164	5.47	1.54
(%)	(2.8)	(4.4)	(7.1)	(11.6)	(13.5)	(15.0)	(12.6)	(1.7)	(7.6)	(8.4)	(5.0)	(3.9)	(2.3)	(1.5)	(2.6)	(100.0)				
その他	264	306	470	727	751	782	606	105	359	317	200	130	60	45	99	5,221	185	5,406	5.12	1.49
(%)	(5.1)	(5.9)	(9.0)	(13.9)	(14.4)	(15.0)	(11.6)	(2.0)	(6.9)	(6.1)	(3.8)	(2.5)	(1.1)	(0.9)	(1.9)	(100.0)				
合計	9,125	10,311	18,021	25,463	30,402	30,654	25,842	4,390	15,034	13,045	7,976	4,862	2,860	1,809	2,898	202,692	6,844	209,536	5.17	1.44
(%)	(4.5)	(5.1)	(8.9)	(12.6)	(15.0)	(15.1)	(12.7)	(2.2)	(7.4)	(6.4)	(3.9)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	(%)																			
総計	9,125	10,311	18,021	25,463	30,402	30,654	25,842	4,390	15,034	13,045	7,976	4,862	2,860	1,809	2,898	202,692	6,844	209,536	5.17	1.44
(%)	(4.5)	(5.1)	(8.9)	(12.6)	(15.0)	(15.1)	(12.7)	(2.2)	(7.4)	(6.4)	(3.9)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4123##

(C)Japanese Society for Dialysis Therapy

2016年調査