

表 1164 透析前リン濃度(mg/dL) 透析歴(年)別 (血液透析濾過患者 透析回数週3回)

透析歴	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
2年未満	355	510	904	1,313	1,465	1,389	1,192	191	661	657	424	266	179	100	166	9,772	246	10,018	5.22	1.46
(%)	(3.6)	(5.2)	(9.3)	(13.4)	(15.0)	(14.2)	(12.2)	(2.0)	(6.8)	(6.7)	(4.3)	(2.7)	(1.8)	(1.0)	(1.7)	(100.0)				
2年～	526	697	1,263	1,874	2,307	2,341	2,111	332	1,219	1,074	705	447	265	150	257	15,568	370	15,938	5.30	1.44
(%)	(3.4)	(4.5)	(8.1)	(12.0)	(14.8)	(15.0)	(13.6)	(2.1)	(7.8)	(6.9)	(4.5)	(2.9)	(1.7)	(1.0)	(1.7)	(100.0)				
5年～	561	672	1,237	1,904	2,538	2,549	2,364	408	1,397	1,289	822	504	356	218	321	17,140	442	17,582	5.39	1.46
(%)	(3.3)	(3.9)	(7.2)	(11.1)	(14.8)	(14.9)	(13.8)	(2.4)	(8.2)	(7.5)	(4.8)	(2.9)	(2.1)	(1.3)	(1.9)	(100.0)				
10年～	301	391	715	1,126	1,429	1,522	1,356	257	796	755	456	285	165	122	172	9,848	268	10,116	5.37	1.44
(%)	(3.1)	(4.0)	(7.3)	(11.4)	(14.5)	(15.5)	(13.8)	(2.6)	(8.1)	(7.7)	(4.6)	(2.9)	(1.7)	(1.2)	(1.7)	(100.0)				
15年～	149	213	399	659	872	931	923	147	536	462	273	128	91	55	86	5,924	134	6,058	5.38	1.34
(%)	(2.5)	(3.6)	(6.7)	(11.1)	(14.7)	(15.7)	(15.6)	(2.5)	(9.0)	(7.8)	(4.6)	(2.2)	(1.5)	(0.9)	(1.5)	(100.0)				
20年～	117	122	288	454	616	646	548	106	296	232	157	92	49	29	30	3,782	75	3,857	5.25	1.30
(%)	(3.1)	(3.2)	(7.6)	(12.0)	(16.3)	(17.1)	(14.5)	(2.8)	(7.8)	(6.1)	(4.2)	(2.4)	(1.3)	(0.8)	(0.8)	(100.0)				
25年～	57	98	179	309	358	375	330	57	167	152	86	61	26	19	16	2,290	55	2,345	5.22	1.29
(%)	(2.5)	(4.3)	(7.8)	(13.5)	(15.6)	(16.4)	(14.4)	(2.5)	(7.3)	(6.6)	(3.8)	(2.7)	(1.1)	(0.8)	(0.7)	(100.0)				
30年～	35	54	81	156	180	234	224	39	109	105	62	27	11	6	5	1,328	39	1,367	5.28	1.21
(%)	(2.6)	(4.1)	(6.1)	(11.7)	(13.6)	(17.6)	(16.9)	(2.9)	(8.2)	(7.9)	(4.7)	(2.0)	(0.8)	(0.5)	(0.4)	(100.0)				
35年～	30	37	59	87	128	123	118	19	70	36	28	13	7	4	5	764	16	780	5.16	1.35
(%)	(3.9)	(4.8)	(7.7)	(11.4)	(16.8)	(16.1)	(15.4)	(2.5)	(9.2)	(4.7)	(3.7)	(1.7)	(0.9)	(0.5)	(0.7)	(100.0)				
40年～	11	15	26	30	43	56	37	5	22	15	8	8	1	2	1	280	7	287	5.08	1.27
(%)	(3.9)	(5.4)	(9.3)	(10.7)	(15.4)	(20.0)	(13.2)	(1.8)	(7.9)	(5.4)	(2.9)	(2.9)	(0.4)	(0.7)	(0.4)	(100.0)				
合計	2,142	2,809	5,151	7,912	9,936	10,166	9,203	1,561	5,273	4,777	3,021	1,831	1,150	705	1,059	66,696	1,652	68,348	5.32	1.42
(%)	(3.2)	(4.2)	(7.7)	(11.9)	(14.9)	(15.2)	(13.8)	(2.3)	(7.9)	(7.2)	(4.5)	(2.7)	(1.7)	(1.1)	(1.6)	(100.0)				
不明	1	1	2	3	3	2	5		1	1	1				1	21	5	26	5.24	1.59
(%)	(4.8)	(4.8)	(9.5)	(14.3)	(14.3)	(9.5)	(23.8)		(4.8)	(4.8)	(4.8)				(4.8)	(100.0)				
記載なし																				
(%)																				
総計	2,143	2,810	5,153	7,915	9,939	10,168	9,208	1,561	5,274	4,778	3,022	1,831	1,150	705	1,060	66,717	1,657	68,374	5.32	1.42
(%)	(3.2)	(4.2)	(7.7)	(11.9)	(14.9)	(15.2)	(13.8)	(2.3)	(7.9)	(7.2)	(4.5)	(2.7)	(1.7)	(1.1)	(1.6)	(100.0)				
平均	8.56	8.60	8.68	9.01	9.22	9.55	9.59	9.81	9.44	9.08	8.94	8.57	7.98	8.38	7.48	9.14	8.96	9.14		
標準偏差	8.43	8.63	8.52	8.59	8.58	8.75	8.66	8.55	8.47	8.11	8.10	7.93	7.20	7.24	6.55	8.48	8.36	8.47		

数値下のかっこ内は行方向の合計に対する％です。  
##D4132##  
(C)Japanese Society for Dialysis Therapy  
2016年調査