

表 1183 PTH値(pg/mL) 原疾患別 (施設血液透析患者 透析回数週3回) PTH測定法:intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,263	3,087	3,309	3,582	3,749	3,640	3,590	3,309	3,024	7,025	81	4,252	2,364	1,417	887	236	344	445	762	616	47,982	818	48,800	185.54	189.64
(%)	(4.7)	(6.4)	(6.9)	(7.5)	(7.8)	(7.6)	(7.5)	(6.9)	(6.3)	(14.6)	(0.2)	(8.9)	(4.9)	(3.0)	(1.8)	(0.5)	(0.7)	(0.9)	(1.6)	(1.3)	(100.0)				
慢性腎盂腎炎	78	92	117	128	115	123	113	84	86	202	5	132	78	42	35	10	12	18	18	23	1,511	36	1,547	183.29	171.51
(%)	(5.2)	(6.1)	(7.7)	(8.5)	(7.6)	(8.1)	(7.5)	(5.6)	(5.7)	(13.4)	(0.3)	(8.7)	(5.2)	(2.8)	(2.3)	(0.7)	(0.8)	(1.2)	(1.2)	(1.5)	(100.0)				
急速進行性糸球体腎炎	48	82	113	135	109	113	109	90	99	205	3	114	46	29	20	2	6	7	8	11	1,349	29	1,378	164.21	156.47
(%)	(3.6)	(6.1)	(8.4)	(10.0)	(8.1)	(8.4)	(8.1)	(6.7)	(7.3)	(15.2)	(0.2)	(8.5)	(3.4)	(2.1)	(1.5)	(0.1)	(0.4)	(0.5)	(0.6)	(0.8)	(100.0)				
妊娠腎/妊娠中毒症	42	32	49	59	52	58	69	43	36	99	1	54	42	16	15	1	10	12	15	11	716	8	724	198.24	237.95
(%)	(5.9)	(4.5)	(6.8)	(8.2)	(7.3)	(8.1)	(9.6)	(6.0)	(5.0)	(13.8)	(0.1)	(7.5)	(5.9)	(2.2)	(2.1)	(0.1)	(1.4)	(1.7)	(2.1)	(1.5)	(100.0)				
その他の分類不能の腎炎	34	48	37	58	68	66	61	56	51	105	3	63	46	24	13	3	4	6	15	9	770	7	777	192.88	248.20
(%)	(4.4)	(6.2)	(4.8)	(7.5)	(8.8)	(8.6)	(7.9)	(7.3)	(6.6)	(13.6)	(0.4)	(8.2)	(6.0)	(3.1)	(1.7)	(0.4)	(0.5)	(0.8)	(1.9)	(1.2)	(100.0)				
多発性嚢胞腎	233	381	435	479	475	460	514	483	411	915	11	549	323	178	123	30	47	67	75	77	6,266	136	6,402	185.16	184.19
(%)	(3.7)	(6.1)	(6.9)	(7.6)	(7.6)	(7.3)	(8.2)	(7.7)	(6.6)	(14.6)	(0.2)	(8.8)	(7.3)	(2.8)	(2.0)	(0.6)	(0.8)	(1.1)	(1.2)	(1.0)	(100.0)				
腎硬化症	631	1,162	1,292	1,418	1,483	1,458	1,376	1,244	1,196	2,717	33	1,574	863	444	287	80	117	116	194	161	17,846	408	18,254	175.11	161.66
(%)	(3.5)	(6.5)	(7.2)	(7.9)	(8.3)	(8.2)	(7.7)	(7.0)	(6.7)	(15.2)	(0.2)	(8.8)	(4.8)	(2.5)	(1.6)	(0.4)	(0.7)	(0.7)	(1.1)	(0.9)	(100.0)				
悪性高血圧	50	73	100	99	112	124	103	106	108	204	5	114	80	51	24	3	11	9	19	27	1,422	35	1,457	189.50	177.34
(%)	(3.5)	(5.1)	(7.0)	(7.0)	(7.9)	(8.7)	(7.2)	(7.5)	(7.6)	(14.3)	(0.4)	(8.0)	(5.6)	(3.6)	(1.7)	(0.2)	(0.8)	(0.6)	(1.3)	(1.9)	(100.0)				
糖尿病性腎症	3,060	4,921	5,334	5,494	5,798	5,459	5,506	5,208	4,464	10,350	141	5,998	3,351	1,848	1,059	257	396	421	671	420	70,156	1,704	71,860	167.58	145.27
(%)	(4.4)	(7.0)	(7.6)	(7.8)	(8.3)	(7.8)	(7.8)	(7.4)	(6.4)	(14.8)	(0.2)	(8.5)	(4.8)	(2.6)	(1.5)	(0.4)	(0.6)	(0.6)	(1.0)	(0.6)	(100.0)				
SLE腎炎	73	88	104	87	79	90	90	89	50	150	2	82	55	28	10	8	7	11	15	12	1,130	19	1,149	166.08	157.99
(%)	(6.5)	(7.8)	(9.2)	(7.7)	(7.0)	(8.0)	(8.0)	(7.9)	(4.4)	(13.3)	(0.2)	(7.3)	(4.9)	(2.5)	(0.9)	(0.7)	(0.6)	(1.0)	(1.3)	(1.1)	(100.0)				
アミロイド腎	9	21	16	21	16	20	19	22	13	28		13	14	3	1	1	1	1	2	2	223	4	227	159.00	152.69
(%)	(4.0)	(9.4)	(7.2)	(9.4)	(7.2)	(9.0)	(8.5)	(9.9)	(5.8)	(12.6)		(5.8)	(6.3)	(1.3)	(0.4)	(0.4)	(0.4)	(0.4)	(0.9)	(0.9)	(100.0)				
痛風腎	26	31	42	51	46	52	46	42	37	72		50	37	22	14	3	3	6	4	10	594	12	606	193.71	250.72
(%)	(4.4)	(5.2)	(7.1)	(8.6)	(7.7)	(8.8)	(7.7)	(7.1)	(6.2)	(12.1)		(8.4)	(6.2)	(3.7)	(2.4)	(0.5)	(0.5)	(1.0)	(0.7)	(1.7)	(100.0)				
先天性代謝異常による腎不全	9	9	7	8	11	10	10	12	5	17		8	9	3	6	2	2	4	2		134	4	138	188.68	153.85
(%)	(6.7)	(6.7)	(5.2)	(6.0)	(8.2)	(7.5)	(7.5)	(9.0)	(3.7)	(12.7)		(6.0)	(6.7)	(2.2)	(4.5)	(1.5)	(1.5)	(3.0)	(1.5)		(100.0)				
腎・尿路結核	8	8	7	5	6	11	5	4	8	13		6	6	1		1					89	1	90	139.11	100.79
(%)	(9.0)	(9.0)	(7.9)	(5.6)	(6.7)	(12.4)	(5.6)	(4.5)	(9.0)	(14.6)		(6.7)	(6.7)	(1.1)		(1.1)					(100.0)				
腎・尿路結石	15	21	21	24	29	32	29	20	16	54		32	14	9	1		4		6	2	329	11	340	168.78	150.05
(%)	(4.6)	(6.4)	(6.4)	(7.3)	(8.8)	(9.7)	(8.8)	(6.1)	(4.9)	(16.4)		(9.7)	(4.3)	(2.7)	(0.3)		(1.2)		(1.8)	(0.6)	(100.0)				
腎・尿路腫瘍	19	25	31	45	53	46	44	36	26	77	3	51	24	17	13	2	9	3	13	4	541	11	552	187.63	160.09
(%)	(3.5)	(4.6)	(5.7)	(8.3)	(9.8)	(8.5)	(8.1)	(6.7)	(4.8)	(14.2)	(0.6)	(9.4)	(4.4)	(3.1)	(2.4)	(0.4)	(1.7)	(0.6)	(2.4)	(0.7)	(100.0)				
閉塞性尿路障害	14	24	25	29	27	31	33	20	23	61		28	24	15	4		3	3	8	8	380	9	389	201.75	248.98
(%)	(3.7)	(6.3)	(6.6)	(7.6)	(7.1)	(8.2)	(8.7)	(5.3)	(6.1)	(16.1)		(7.4)	(6.3)	(3.9)	(1.1)		(0.8)	(0.8)	(2.1)	(2.1)	(100.0)				
骨髄腫	9	12	16	13	6	13	16	15	4	37		10	7	3	2	2		1	1		167	3	170	155.86	115.57
(%)	(5.4)	(7.2)	(9.6)	(7.8)	(3.6)	(7.8)	(9.6)	(9.0)	(2.4)	(22.2)		(6.0)	(4.2)	(1.8)	(1.2)	(1.2)		(0.6)	(0.6)		(100.0)				
腎形成不全	10	24	21	14	26	19	19	19	19	55		36	16	10	7	2	2	1	6	6	312	4	316	203.54	210.91
(%)	(3.2)	(7.7)	(6.7)	(4.5)	(8.3)	(6.1)	(6.1)	(6.1)	(6.1)	(17.6)		(11.5)	(5.1)	(3.2)	(2.2)	(0.6)	(0.6)	(0.3)	(1.9)	(1.9)	(100.0)				
不明	662	1,104	1,273	1,366	1,406	1,312	1,341	1,214	1,037	2,505	40	1,448	802	512	306	68	113	142	211	205	17,067	391	17,458	181.29	190.29
(%)	(3.9)	(6.5)	(7.5)	(8.0)	(8.2)	(7.7)	(7.9)	(7.1)	(6.1)	(14.7)	(0.2)	(8.5)	(4.7)	(3.0)	(1.8)	(0.4)	(0.7)	(0.8)	(1.2)	(1.2)	(100.0)				
再導入	58	74	56	77	74	57	66	61	67	107	1	99	55	33	18	4	6	9	16	18	956	10	966	191.06	197.80
(%)	(6.1)	(7.7)	(5.9)	(8.1)	(7.7)	(6.0)	(6.9)	(6.4)	(7.0)	(11.2)	(0.1)	(10.4)	(5.8)	(3.5)	(1.9)	(0.4)	(0.6)	(0.9)	(1.7)	(1.9)	(100.0)				
その他	187	293	333	354	338	359	350	302	311	635	11	390	196	118	91	27	31	28	60	61	4,475	128	4,603	181.51	179.82
(%)	(4.2)	(6.5)	(7.4)	(7.9)	(7.6)	(8.0)	(7.8)	(6.7)	(6.9)	(14.2)	(0.2)	(8.7)	(4.4)	(2.6)	(2.0)	(0.6)	(0.7)	(0.6)	(1.3)	(1.4)	(100.0)				
合計	7,538	11,612	12,738	13,546	14,078	13,553	13,509	12,479	11,091	25,633	340	15,103	8,452	4,823	2,936	742	1,128	1,310	2,121	1,683	174,415	3,788	178,203	176.54	170.05
(%)	(4.3)	(6.7)	(7.3)	(7.8)	(8.1)	(7.8)	(7.7)	(7.2)	(6.4)	(14.7)	(0.2)	(8.7)	(4.8)	(2.8)	(1.7)	(0.4)	(0.6)	(0.8)	(1.2)	(1.0)	(100.0)				
記載なし																									
(%)																									
総計	7,538	11,612	12,738	13,546	14,078	13,553	13,509	12,479	11,091	25,633	340	15,103	8,452	4,823	2,936	742	1,128	1,310	2,121	1,683	174,415	3,788	178,203	176.54	170.05
(%)	(4.3)	(6.7)	(7.3)	(7.8)	(8.1)	(7.8)	(7.7)	(7.2)	(6.4)	(14.7)	(0.2)	(8.7)	(4.8)	(2.8)	(1.7)	(0.4)	(0.6)	(0.8)	(1.2)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。  
##D4223##  
(C)Japanese Society for Dialysis Therapy  
2016年調査