

表 1188 PTH値(pg/mL) 原疾患別(血液透析濾過患者 透析回数週3回) PTH測定法:intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	883	1,070	1,300	1,364	1,529	1,537	1,498	1,447	1,250	2,888	39	1,734	1,007	551	374	84	148	136	289	271	19,399	243	19,642	187.42	192.05
(%)	(4.6)	(5.5)	(6.7)	(7.0)	(7.9)	(7.9)	(7.7)	(7.5)	(6.4)	(14.9)	(0.2)	(8.9)	(5.2)	(2.8)	(1.9)	(0.4)	(0.8)	(0.7)	(1.5)	(1.4)	(100.0)				
慢性腎盂腎炎	32	23	40	43	49	42	36	35	41	78		38	32	17	11	1	4	6	6	4	538	12	550	179.03	179.44
(%)	(5.9)	(4.3)	(7.4)	(8.0)	(9.1)	(7.8)	(6.7)	(6.5)	(7.6)	(14.5)		(7.1)	(5.9)	(3.2)	(2.0)	(0.2)	(0.7)	(1.1)	(1.1)	(0.7)	(100.0)				
急速進行性糸球体腎炎	19	34	24	34	28	32	34	32	29	61		38	16	16	9	1	4	2	6	4	423	4	427	180.44	172.82
(%)	(4.5)	(8.0)	(5.7)	(8.0)	(6.6)	(7.6)	(8.0)	(7.6)	(6.9)	(14.4)		(9.0)	(3.8)	(3.8)	(2.1)	(0.2)	(0.9)	(0.5)	(1.4)	(0.9)	(100.0)				
妊娠腎/妊娠中毒症	26	27	30	25	33	24	37	24	31	50	3	26	25	10	5	1	4	1	5	8	395	3	398	178.46	172.58
(%)	(6.6)	(6.8)	(7.6)	(6.3)	(8.4)	(6.1)	(9.4)	(6.1)	(7.8)	(12.7)	(0.8)	(6.6)	(6.3)	(2.5)	(1.3)	(0.3)	(1.0)	(0.3)	(1.3)	(2.0)	(100.0)				
その他の分類不能の腎炎	12	17	12	19	20	25	23	17	25	51		35	16	12	5	3	4	4	6	5	311	9	320	209.17	194.17
(%)	(3.9)	(5.5)	(3.9)	(6.1)	(6.4)	(8.0)	(7.4)	(5.5)	(8.0)	(16.4)		(11.3)	(5.1)	(3.9)	(1.6)	(1.0)	(1.3)	(1.3)	(1.9)	(1.6)	(100.0)				
多発性囊胞腎	75	119	142	163	184	184	168	141	151	371	5	193	108	61	42	8	20	17	34	24	2,210	33	2,243	186.24	178.36
(%)	(3.4)	(5.4)	(6.4)	(7.4)	(8.3)	(8.3)	(7.6)	(6.4)	(6.8)	(16.8)	(0.2)	(8.7)	(4.9)	(2.8)	(1.9)	(0.4)	(0.9)	(0.8)	(1.5)	(1.1)	(100.0)				
腎硬化症	159	304	348	428	443	395	408	380	324	761	10	458	260	150	95	24	34	41	47	41	5,110	72	5,182	177.41	158.47
(%)	(3.1)	(5.9)	(6.8)	(8.4)	(8.7)	(7.7)	(8.0)	(7.4)	(6.3)	(14.9)	(0.2)	(9.0)	(5.1)	(2.9)	(1.9)	(0.5)	(0.7)	(0.8)	(0.9)	(0.8)	(100.0)				
悪性高血圧	11	20	35	42	43	48	32	32	29	76	1	42	24	15	9		4	4	7	5	479	4	483	186.94	181.18
(%)	(2.3)	(4.2)	(7.3)	(8.8)	(9.0)	(10.0)	(6.7)	(6.7)	(6.1)	(15.9)	(0.2)	(8.8)	(5.0)	(3.1)	(1.9)		(0.8)	(0.8)	(1.5)	(1.0)	(100.0)				
糖尿病性腎症	786	1,380	1,526	1,688	1,773	1,736	1,727	1,507	1,515	3,158	35	1,812	1,001	545	321	68	117	140	200	121	21,156	309	21,465	168.28	138.12
(%)	(3.7)	(6.5)	(7.2)	(8.0)	(8.4)	(8.2)	(8.2)	(7.1)	(7.2)	(14.9)	(0.2)	(8.6)	(4.7)	(2.6)	(1.5)	(0.3)	(0.6)	(0.7)	(0.9)	(0.6)	(100.0)				
SLE腎炎	22	34	41	47	24	30	37	24	36	55	1	37	28	14	8	2	3	4	5	2	454	8	462	168.41	141.08
(%)	(4.8)	(7.5)	(9.0)	(10.4)	(5.3)	(6.6)	(8.1)	(5.3)	(7.9)	(12.1)	(0.2)	(8.1)	(6.2)	(3.1)	(1.8)	(0.4)	(0.7)	(0.9)	(1.1)	(0.4)	(100.0)				
アミロイド腎	5	9	5	5	7	5	2	7	5	14		6	4		2			2	1	2	81	3	84	180.40	183.64
(%)	(6.2)	(11.1)	(6.2)	(6.2)	(8.6)	(6.2)	(2.5)	(8.6)	(6.2)	(17.3)		(7.4)	(4.9)		(2.5)			(2.5)	(1.2)	(2.5)	(100.0)				
痛風腎	12	6	11	16	11	12	12	16	16	27		15	10	5	9	5		2	7	3	197	3	200	209.43	176.49
(%)	(6.1)	(3.0)	(5.6)	(8.1)	(5.6)	(6.1)	(6.1)	(9.1)	(8.1)	(13.7)		(7.6)	(5.1)	(2.5)	(4.6)	(2.5)		(1.0)	(3.6)	(1.5)	(100.0)				
先天性代謝異常による腎不全	4	3	3	7	3	4	3	4	5	9	1	8	3	3					4		64		64	195.02	158.74
(%)	(6.3)	(4.7)	(4.7)	(10.9)	(4.7)	(6.3)	(4.7)	(6.3)	(7.8)	(14.1)	(1.6)	(12.5)	(4.7)	(4.7)					(6.3)		(100.0)				
腎・尿路結核	3	3	2	4	1	2	4	1	1	5		2		4	1						33	2	35	158.58	132.04
(%)	(9.1)	(9.1)	(6.1)	(12.1)	(3.0)	(6.1)	(12.1)	(3.0)	(3.0)	(15.2)		(6.1)		(12.1)	(3.0)						(100.0)				
腎・尿路結石	2	8	8	3	7	9	5	3	6	13		10	6	2	2	2		1	1	1	89	2	91	192.52	179.25
(%)	(2.2)	(9.0)	(9.0)	(3.4)	(7.9)	(10.1)	(5.6)	(3.4)	(6.7)	(14.6)		(11.2)	(6.7)	(2.2)	(2.2)	(2.2)		(1.1)	(1.1)	(1.1)	(100.0)				
腎・尿路腫瘍	7	9	11	10	16	16	11	9	16	29		7	12	3	2	1	1	3	1		164	2	166	168.07	122.36
(%)	(4.3)	(5.5)	(6.7)	(6.1)	(9.8)	(9.8)	(6.7)	(5.5)	(9.8)	(17.7)		(4.3)	(7.3)	(1.8)	(1.2)	(0.6)	(0.6)	(1.8)	(0.6)		(100.0)				
閉塞性尿路障害	11	7	8	13	8	7	12	5	5	22		15	7	5	3	2		1	1	2	134		134	180.54	150.60
(%)	(8.2)	(5.2)	(6.0)	(9.7)	(6.0)	(5.2)	(9.0)	(3.7)	(3.7)	(16.4)		(11.2)	(5.2)	(3.7)	(2.2)	(1.5)		(0.7)	(0.7)	(1.5)	(100.0)				
骨髄腫	3	1	1	2	2	2	2	1	3	7	1	4	1						1		29	2	31	183.52	136.15
(%)		(10.3)	(3.4)	(3.4)	(6.9)	(6.9)	(6.9)	(3.4)	(10.3)	(24.1)	(3.4)	(13.8)	(3.4)						(3.4)		(100.0)				
腎形成不全	4	8	11	4	7	8	14	9	8	11	1	24	12	4	4	1	4	1	2	5	142	3	145	237.42	237.19
(%)	(2.8)	(5.6)	(7.7)	(2.8)	(4.9)	(5.6)	(9.9)	(6.3)	(5.6)	(7.7)	(0.7)	(16.9)	(8.5)	(2.8)	(2.8)	(0.7)	(2.8)	(0.7)	(1.4)	(3.5)	(100.0)				
不明	232	320	372	461	464	455	450	371	358	806	10	520	263	178	102	21	53	45	86	75	5,642	85	5,727	186.35	183.27
(%)	(4.1)	(5.7)	(6.6)	(8.2)	(8.1)	(8.0)	(6.6)	(6.3)	(14.3)	(0.2)	(9.2)	(4.7)	(3.2)	(1.8)	(0.4)	(0.9)	(0.8)	(1.5)	(1.3)	(100.0)					
再導入	26	29	33	41	42	49	35	38	33	77		44	28	12	11	7	3	3	10	18	539	7	546	201.63	204.16
(%)	(4.8)	(5.4)	(6.1)	(7.6)	(7.8)	(9.1)	(6.5)	(7.1)	(6.1)	(14.3)		(8.2)	(5.2)	(2.2)	(2.0)	(1.3)	(0.6)	(0.6)	(1.9)	(3.3)	(100.0)				
その他	59	112	86	93	126	110	115	106	81	207	2	134	78	49	28	6	11	14	21	22	1,460	27	1,487	187.47	189.73
(%)	(4.0)	(7.7)	(5.9)	(6.4)	(8.6)	(7.5)	(7.9)	(7.3)	(5.5)	(14.2)	(0.1)	(9.2)	(5.3)	(3.4)	(1.9)	(0.4)	(0.8)	(1.0)	(1.4)	(1.5)	(100.0)				
合計	2,390	3,545	4,049	4,511	4,820	4,732	4,665	4,211	3,968	8,776	109	5,202	2,941	1,656	1,043	237	414	427	740	613	59,049	833	59,882	179.57	169.35
(%)	(4.0)	(6.0)	(6.9)	(7.6)	(8.2)	(8.0)	(7.9)	(7.1)	(6.7)	(14.9)	(0.2)	(8.8)	(5.0)	(2.8)	(1.8)	(0.4)	(0.7)	(0.7)	(1.3)	(1.0)	(100.0)				
記載なし																									
(%)																									
総計	2,390	3,545	4,049	4,511	4,820	4,732	4,665	4,211	3,968	8,776	109	5,202	2,941	1,656	1,043	237	414	427	740	613	59,049	833	59,882	179.57	169.35
(%)	(4.0)	(6.0)	(6.9)	(7.6)	(8.2)	(8.0)	(7.9)	(7.1)	(6.7)	(14.9)	(0.2)	(8.8)	(5.0)	(2.8)	(1.8)	(0.4)	(0.7)	(0.7)	(1.3)	(1.0)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

##D4233##

(C)Japanese Society for Dialysis Therapy

2016年調査