

表 1193 PTH値(pg/mL) 原疾患別 (腹膜透析患者) PTH測定法:intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	35	76	96	92	98	103	89	89	82	192	6	139	95	77	38	9	20	14	48	38	1,436	84	1,520	231.98	249.25
(%)	(2.4)	(5.3)	(6.7)	(6.4)	(6.8)	(7.2)	(6.2)	(6.2)	(5.7)	(13.4)	(0.4)	(9.7)	(6.6)	(5.4)	(2.6)	(0.6)	(1.4)	(1.0)	(3.3)	(2.6)	(100.0)				
慢性腎盂腎炎	1	3	5	5	7	3	1	3	2	9		2	6	1	2				2	3	55	4	59	229.11	248.11
(%)	(1.8)	(5.5)	(9.1)	(9.1)	(12.7)	(5.5)	(1.8)	(5.5)	(3.6)	(16.4)		(3.6)	(10.9)	(1.8)	(3.6)				(3.6)	(5.5)	(100.0)				
急速進行性糸球体腎炎		4	2	4	3	2	3	2	2	8		4		1					1	1	37	2	39	192.51	208.98
(%)		(10.8)	(5.4)	(10.8)	(8.1)	(5.4)	(8.1)	(5.4)	(5.4)	(21.6)		(10.8)		(2.7)					(2.7)	(2.7)	(100.0)				
妊娠腎/妊娠中毒症		1			1	1			1	4		2		1			1	1		1	14	3	17	307.21	252.56
(%)		(7.1)			(7.1)	(7.1)			(7.1)	(28.6)		(14.3)		(7.1)			(7.1)	(7.1)		(7.1)	(100.0)				
その他の分類不能の腎炎	1	4	1	2	5	2	3		2	7		3	7	2				1	3	2	45	2	47	273.00	319.87
(%)	(2.2)	(8.9)	(2.2)	(4.4)	(11.1)	(4.4)	(6.7)		(4.4)	(15.6)		(6.7)	(15.6)	(4.4)				(2.2)	(6.7)	(4.4)	(100.0)				
多発性嚢胞腎	5	4	7	10	7	10	9	5	11	28		21	7	4	7	2		4	1	4	146	11	157	244.38	287.83
(%)	(3.4)	(2.7)	(4.8)	(6.8)	(4.8)	(6.8)	(6.2)	(3.4)	(7.5)	(19.2)		(14.4)	(4.8)	(2.7)	(4.8)	(1.4)		(2.7)	(0.7)	(2.7)	(100.0)				
腎硬化症	13	25	42	36	39	41	49	36	28	100	1	57	37	32	11	3	9	7	10	12	588	43	631	212.28	186.29
(%)	(2.2)	(4.3)	(7.1)	(6.1)	(6.6)	(7.0)	(8.3)	(6.1)	(4.8)	(17.0)	(0.2)	(9.7)	(6.3)	(5.4)	(1.9)	(0.5)	(1.5)	(1.2)	(1.7)	(2.0)	(100.0)				
悪性高血圧			5	1	3	1	4	5	4	9		8	5	2	1		1	1	3		53	2	55	237.19	152.99
(%)			(9.4)	(1.9)	(5.7)	(1.9)	(7.5)	(9.4)	(7.5)	(17.0)		(15.1)	(9.4)	(3.8)	(1.9)		(1.9)	(1.9)	(5.7)		(100.0)				
糖尿病性腎症	37	72	84	81	81	90	87	88	63	206		157	88	62	38	10	17	15	35	24	1,335	103	1,438	218.61	192.73
(%)	(2.8)	(5.4)	(6.3)	(6.1)	(6.1)	(6.7)	(6.5)	(6.6)	(4.7)	(15.4)		(11.8)	(6.6)	(4.6)	(2.8)	(0.7)	(1.3)	(1.1)	(2.6)	(1.8)	(100.0)				
SLE腎炎		1	3	2	1	2	3	1		6			2	1	1				1		24		24	197.13	161.36
(%)		(4.2)	(12.5)	(8.3)	(4.2)	(8.3)	(12.5)	(4.2)		(25.0)			(8.3)	(4.2)	(4.2)				(4.2)		(100.0)				
アミロイド腎	1				1			3	1				1	1							8		8	177.75	117.95
(%)	(12.5)				(12.5)			(37.5)	(12.5)				(12.5)	(12.5)							(100.0)				
痛風腎	1	2			1		2	2	1	3		1									13	1	14	141.54	93.64
(%)	(7.7)	(15.4)			(7.7)		(15.4)	(15.4)	(7.7)	(23.1)		(7.7)									(100.0)				
先天性代謝異常による腎不全			2			2	2	1	1				1	2							11		11	185.09	132.85
(%)			(18.2)			(18.2)	(18.2)	(9.1)	(9.1)				(9.1)	(18.2)							(100.0)				
腎・尿路結核																									
(%)																									
腎・尿路結石		1				1	2	1	1			3		1					1		11		11	244.82	209.65
(%)		(9.1)				(9.1)	(18.2)	(9.1)	(9.1)			(27.3)		(9.1)					(9.1)		(100.0)				
腎・尿路腫瘍			1		1							2	1			1					6		6	249.50	162.90
(%)			(16.7)		(16.7)							(33.3)	(16.7)			(16.7)					(100.0)				
閉塞性尿路障害					3	1	1		1	2			1						1		10		10	205.20	168.23
(%)					(30.0)	(10.0)	(10.0)		(10.0)	(20.0)			(10.0)						(10.0)		(100.0)				
骨髄腫	1	2																			3		3	38.33	20.21
(%)	(33.3)		(66.7)																		(100.0)				
腎形成不全	2	5	4	2	2	3	1		1	3	1	3		1	1			2	4	4	39		39	319.85	366.43
(%)	(5.1)	(12.8)	(10.3)	(5.1)	(5.1)	(7.7)	(2.6)		(2.6)	(7.7)	(2.6)	(7.7)		(2.6)	(2.6)			(5.1)	(10.3)	(10.3)	(100.0)				
不明	15	15	24	19	29	23	23	16	31	67	3	64	33	23	12		6	7	10	11	431	29	460	232.51	192.01
(%)	(3.5)	(3.5)	(5.6)	(4.4)	(6.7)	(5.3)	(5.3)	(3.7)	(7.2)	(15.5)	(0.7)	(14.8)	(7.7)	(5.3)	(2.8)		(1.4)	(1.6)	(2.3)	(2.6)	(100.0)				
再導入	2	3	2		1	3	1	3	4	8		3	1		3				2	1	37	3	40	226.73	211.93
(%)	(5.4)	(8.1)	(5.4)		(2.7)	(8.1)	(2.7)	(8.1)	(10.8)	(21.6)		(8.1)	(2.7)		(8.1)				(5.4)	(2.7)	(100.0)				
その他	6	6	6	2	4	17	8	7	6	21	1	9	5	4	6	1	3	1	3	5	121	15	136	252.68	291.76
(%)	(5.0)	(5.0)	(5.0)	(1.7)	(3.3)	(14.0)	(6.6)	(5.8)	(5.0)	(17.4)	(0.8)	(7.4)	(4.1)	(3.3)	(5.0)	(0.8)	(2.5)	(0.8)	(2.5)	(4.1)	(100.0)				
合計	120	222	286	256	287	305	288	262	242	673	12	477	291	215	120	26	57	53	125	106	4,423	302	4,725	226.63	222.22
(%)	(2.7)	(5.0)	(6.5)	(5.8)	(6.5)	(6.9)	(6.5)	(5.9)	(5.5)	(15.2)	(0.3)	(10.8)	(6.6)	(4.9)	(2.7)	(0.6)	(1.3)	(1.2)	(2.8)	(2.4)	(100.0)				
記載なし																									
(%)																									
総計	120	222	286	256	287	305	288	262	242	673	12	477	291	215	120	26	57	53	125	106	4,423	302	4,725	226.63	222.22
(%)	(2.7)	(5.0)	(6.5)	(5.8)	(6.5)	(6.9)	(6.5)	(5.9)	(5.5)	(15.2)	(0.3)	(10.8)	(6.6)	(4.9)	(2.7)	(0.6)	(1.3)	(1.2)	(2.8)	(2.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
##D4243##
(C)Japanese Society for Dialysis Therapy
2016年調査