

表 1196 PTH値(pg/mL) 治療方法(新分類)別 (透析患者全体) PTH測定法:whole-PTH

治療方法 (新分類)	20未満	20~	35~	50~	75~	100~	125~	150~	151~	180~	210~	240~	300~	350~	400~	500~	合計	記載なし	総計	平均	標準偏差	
施設血液透析	1,280	1,695	1,707	2,577	2,212	1,791	1,319	49	968	713	465	539	251	148	181	204	16,099	410	16,509	110.91	116.49	
(%)	(8.0)	(10.5)	(10.6)	(16.0)	(13.7)	(11.1)	(8.2)	(0.3)	(6.0)	(4.4)	(2.9)	(3.3)	(1.6)	(0.9)	(1.1)	(1.3)	(100.0)		268	100.62	98.53	
Off-line HDF	15	39	40	45	40	22	14	1	15	8	8	10	4	2	3	2	268					
(%)	(5.6)	(14.6)	(14.9)	(16.8)	(14.9)	(8.2)	(5.2)	(0.4)	(5.6)	(3.0)	(3.0)	(3.7)	(1.5)	(0.7)	(1.1)	(0.7)	(100.0)		32	4,352	117.85	125.90
On-line HDF	324	411	396	700	621	495	358	13	257	206	129	159	81	47	57	66	4,320					
(%)	(7.5)	(9.5)	(9.2)	(16.2)	(14.4)	(11.5)	(8.3)	(0.3)	(5.9)	(4.8)	(3.0)	(3.7)	(1.9)	(1.1)	(1.3)	(1.5)	(100.0)					
Push/Pull HDF			2	2		4	1		1	2							12		12	111.17	54.28	
(%)			(16.7)	(16.7)		(33.3)	(8.3)		(8.3)	(16.7)							(100.0)					
ハイオフィルトレーション	3	3	5	2	3	3	3		1	2	1	2					28	13	41	96.29	76.00	
(%)	(10.7)	(10.7)	(17.9)	(7.1)	(10.7)	(10.7)	(10.7)		(3.6)	(7.1)	(3.6)	(7.1)					(100.0)					
IHDF	58	66	68	91	83	78	57	2	39	28	25	28	10	9	7	4	653	3	656	110.36	96.51	
(%)	(8.9)	(10.1)	(10.4)	(13.9)	(12.7)	(11.9)	(8.7)	(0.3)	(6.0)	(4.3)	(3.8)	(4.3)	(1.5)	(1.4)	(1.1)	(0.6)	(100.0)					
血液濾過																						
(%)																						
血液吸着透析 (リクセル等使用)	25	9	11	21	20	15	10		9	3	4	1	2	2			132		132	89.41	74.60	
(%)	(18.9)	(6.8)	(8.3)	(15.9)	(15.2)	(11.4)	(7.6)		(6.8)	(2.3)	(3.0)	(0.8)	(1.5)	(1.5)			(100.0)					
在宅血液透析	3	7	9	12	8	6	6	1	5	1	1	3		1	2	1	66	1	67	116.91	122.28	
(%)	(4.5)	(10.6)	(13.6)	(18.2)	(12.1)	(9.1)	(9.1)	(1.5)	(7.6)	(1.5)	(1.5)	(4.5)		(1.5)	(3.0)	(1.5)	(100.0)					
腹膜透析 (CAPD)	30	28	39	64	49	45	39		20	23	15	16	8	6	6	7	395	27	422	123.57	114.91	
(%)	(7.6)	(7.1)	(9.9)	(16.2)	(12.4)	(11.4)	(9.9)		(5.1)	(5.8)	(3.8)	(4.1)	(2.0)	(1.5)	(1.5)	(1.8)	(100.0)					
腹膜透析 (APD)	4	13	10	13	13	17	10		10	8	4	10	7	2	1	2	124	10	134	146.96	142.97	
(%)	(3.2)	(10.5)	(8.1)	(10.5)	(10.5)	(13.7)	(8.1)		(8.1)	(6.5)	(3.2)	(8.1)	(5.6)	(1.6)	(0.8)	(1.6)	(100.0)					
腹膜透析 (CCPD)	5	7	9	15	8	9	6		9	4	4	4	2	1	2	1	86	1	87	124.90	105.96	
(%)	(5.8)	(8.1)	(10.5)	(17.4)	(9.3)	(10.5)	(7.0)		(10.5)	(4.7)	(4.7)	(4.7)	(2.3)	(1.2)	(1.2)	(1.2)	(100.0)					
合計	1,747	2,278	2,296	3,542	3,057	2,485	1,823	66	1,334	998	656	772	365	218	259	287	22,183	497	22,680	112.47	117.57	
(%)	(7.9)	(10.3)	(10.4)	(16.0)	(13.8)	(11.2)	(8.2)	(0.3)	(6.0)	(4.5)	(3.0)	(3.5)	(1.6)	(1.0)	(1.2)	(1.3)	(100.0)					
記載なし																						
(%)																						
総計	1,747	2,278	2,296	3,542	3,057	2,485	1,823	66	1,334	998	656	772	365	218	259	287	22,183	497	22,680	112.47	117.57	
(%)	(7.9)	(10.3)	(10.4)	(16.0)	(13.8)	(11.2)	(8.2)	(0.3)	(6.0)	(4.5)	(3.0)	(3.5)	(1.6)	(1.0)	(1.2)	(1.3)	(100.0)					

数値下の括弧内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

##D4251##

(C)Japanese Society for Dialysis Therapy

2016年調査