

表 1197 PTH値(pg/mL) 併用の状況別 (透析患者全体) PTH測定法:whole-PTH

併用の状況	20未満	20~	35~	50~	75~	100~	125~	150~	151~	180~	210~	240~	300~	350~	400~	500~	合計	記載なし	総計	平均	標準偏差
HD(F)単独	1,707	2,229	2,237	3,445	2,984	2,413	1,767	66	1,292	963	633	741	348	207	250	276	21,558	459	22,017	111.99	117.44
(%)	(7.9)	(10.3)	(10.4)	(16.0)	(13.8)	(11.2)	(8.2)	(0.3)	(6.0)	(4.5)	(2.9)	(3.4)	(1.6)	(1.0)	(1.2)	(1.3)	(100.0)				
PD単独	35	35	37	60	46	54	45		29	30	20	25	15	8	4	8	451	33	484	133.26	123.64
(%)	(7.8)	(7.8)	(8.2)	(13.3)	(10.2)	(12.0)	(10.0)		(6.4)	(6.7)	(4.4)	(5.5)	(3.3)	(1.8)	(0.9)	(1.8)	(100.0)				
HD(F)単独,PDカテあり				3	3	1	1		2				1				11	11	127.27	91.29	
(%)				(27.3)	(27.3)	(9.1)	(9.1)		(18.2)				(9.1)				(100.0)				
PD,HD併用週1HD(F)	4	9	14	20	14	11	8		5	4	2	3	2	1	5	2	104	5	109	124.28	124.80
(%)	(3.8)	(8.7)	(13.5)	(19.2)	(13.5)	(10.6)	(7.7)		(4.8)	(3.8)	(1.9)	(2.9)	(1.9)	(1.0)	(4.8)	(1.9)	(100.0)				
PD,HD併用週2HD(F)		4	6	13	9	6	2		5	1	1	3					50	50	99.46	64.56	
(%)		(8.0)	(12.0)	(26.0)	(18.0)	(12.0)	(4.0)		(10.0)	(2.0)	(2.0)	(6.0)					(100.0)				
PD,HD併用週3HD(F)	1		1						1				1			1	5	5	229.80	237.38	
(%)	(20.0)		(20.0)						(20.0)				(20.0)			(20.0)	(100.0)				
PD,HD併用週4HD(F)																					
(%)																					
PD,HD併用その他		1	1	1	1											4	4	53.75	25.82		
(%)		(25.0)	(25.0)	(25.0)	(25.0)											(100.0)					
合計	1,747	2,278	2,296	3,542	3,057	2,485	1,823	66	1,334	998	656	772	365	218	259	287	22,183	497	22,680	112.47	117.57
(%)	(7.9)	(10.3)	(10.4)	(16.0)	(13.8)	(11.2)	(8.2)	(0.3)	(6.0)	(4.5)	(3.0)	(3.5)	(1.6)	(1.0)	(1.2)	(1.3)	(100.0)				
不明																					
(%)																					
記載なし																					
(%)																					
総計	1,747	2,278	2,296	3,542	3,057	2,485	1,823	66	1,334	998	656	772	365	218	259	287	22,183	497	22,680	112.47	117.57
(%)	(7.9)	(10.3)	(10.4)	(16.0)	(13.8)	(11.2)	(8.2)	(0.3)	(6.0)	(4.5)	(3.0)	(3.5)	(1.6)	(1.0)	(1.2)	(1.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4252##

(C)Japanese Society for Dialysis Therapy
2016年調査