

表 1211 PTH値(pg/mL) 原疾患別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

原疾患	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	130	183	182	259	219	188	142	6	97	79	63	76	42	21	25	31	1,743	13	1,756	122.34	137.67
(%)	(7.5)	(10.5)	(10.4)	(14.9)	(12.6)	(10.8)	(8.1)	(0.3)	(5.6)	(4.5)	(3.6)	(4.4)	(2.4)	(1.2)	(1.4)	(1.8)	(100.0)				
慢性腎盂腎炎	4	5	4	6	9	7	4	1	2		2	1	1	1		2	49	1	50	124.41	138.42
(%)	(8.2)	(10.2)	(8.2)	(12.2)	(18.4)	(14.3)	(8.2)	(2.0)	(4.1)		(4.1)	(2.0)	(2.0)	(2.0)		(4.1)	(100.0)				
急速進行性糸球体腎炎	3	2	3	8	3	8	1		3	1	1			1	1	1	36		36	121.81	141.32
(%)	(8.3)	(5.6)	(8.3)	(22.2)	(8.3)	(22.2)	(2.8)		(8.3)	(2.8)	(2.8)			(2.8)	(2.8)	(2.8)	(100.0)				
妊娠腎/妊娠中毒症	4	5	2	5	8	1	5		1							1	32		32	123.09	289.92
(%)	(12.5)	(15.6)	(6.3)	(15.6)	(25.0)	(3.1)	(15.6)		(3.1)							(3.1)	(100.0)				
その他の分類不能の腎炎	4	3	1	1	5	4			1	1		1			1		22		22	97.95	99.82
(%)	(18.2)	(13.6)	(4.5)	(4.5)	(22.7)	(18.2)			(4.5)	(4.5)		(4.5)			(4.5)		(100.0)				
多発性嚢胞腎	15	18	25	39	23	21	27		16	11	3	13	5	2	6	1	225	2	227	119.59	106.17
(%)	(6.7)	(8.0)	(11.1)	(17.3)	(10.2)	(9.3)	(12.0)		(7.1)	(4.9)	(1.3)	(5.8)	(2.2)	(0.9)	(2.7)	(0.4)	(100.0)				
腎硬化症	41	37	41	80	69	35	35	1	21	18	8	15	9	4	4	8	426	1	427	111.30	120.64
(%)	(9.6)	(8.7)	(9.6)	(18.8)	(16.2)	(8.2)	(8.2)	(0.2)	(4.9)	(4.2)	(1.9)	(3.5)	(2.1)	(0.9)	(0.9)	(1.9)	(100.0)				
悪性高血圧	2	2	3	3	12	11	4			2		2	1				42	1	43	108.00	65.41
(%)	(4.8)	(4.8)	(7.1)	(7.1)	(28.6)	(26.2)	(9.5)			(4.8)		(4.8)	(2.4)				(100.0)				
糖尿病性腎症	138	170	163	300	260	211	146	3	121	84	57	52	20	15	16	12	1,768	14	1,782	106.53	91.60
(%)	(7.8)	(9.6)	(9.2)	(17.0)	(14.7)	(11.9)	(8.3)	(0.2)	(6.8)	(4.8)	(3.2)	(2.9)	(1.1)	(0.8)	(0.9)	(0.7)	(100.0)				
SLE腎炎	3	8		12	8	6	4			3	2	1	1	1			49	2	51	101.10	81.58
(%)	(6.1)	(16.3)		(24.5)	(16.3)	(12.2)	(8.2)			(6.1)	(4.1)	(2.0)	(2.0)	(2.0)			(100.0)				
アミロイド腎			3	1	1	1	1										7		7	76.00	38.88
(%)			(42.9)	(14.3)	(14.3)	(14.3)	(14.3)										(100.0)				
痛風腎	3	3	1		2	2	2			3		1	1			1	19		19	144.37	166.16
(%)	(15.8)	(15.8)	(5.3)		(10.5)	(10.5)	(10.5)			(15.8)		(5.3)	(5.3)			(5.3)	(100.0)				
先天性代謝異常による腎不全		1	2	1		1				1							6		6	83.83	66.45
(%)		(16.7)	(33.3)	(16.7)		(16.7)				(16.7)							(100.0)				
腎・尿路結核		1	1	1	1	1	1		1								7		7	89.43	49.15
(%)		(14.3)	(14.3)	(14.3)	(14.3)	(14.3)	(14.3)		(14.3)								(100.0)				
腎・尿路結石			1	4	1		2	1		1	1	1	1				13		13	138.38	89.55
(%)			(7.7)	(30.8)	(7.7)		(15.4)	(7.7)		(7.7)	(7.7)	(7.7)	(7.7)				(100.0)				
腎・尿路腫瘍	1	3		1	1	1	1		1	1	1						11		11	94.91	74.87
(%)	(9.1)	(27.3)		(9.1)	(9.1)	(9.1)	(9.1)		(9.1)	(9.1)	(9.1)						(100.0)				
閉塞性尿路障害	3	1		1	1	3	1		1	2				1			14		14	114.21	98.71
(%)	(21.4)	(7.1)		(7.1)	(7.1)	(21.4)	(7.1)		(7.1)	(14.3)				(7.1)			(100.0)				
骨髄腫			1			1									1		3		3	207.00	231.93
(%)			(33.3)			(33.3)									(33.3)		(100.0)				
腎形成不全	1	2	2	4	1	2			1	2	1				1		17		17	108.82	99.81
(%)	(5.9)	(11.8)	(11.8)	(23.5)	(5.9)	(11.8)			(5.9)	(11.8)	(5.9)				(5.9)		(100.0)				
不明	30	42	48	67	74	55	40	2	29	21	13	20	7	6	7	10	471	9	480	123.90	136.91
(%)	(6.4)	(8.9)	(10.2)	(14.2)	(15.7)	(11.7)	(8.5)	(0.4)	(6.2)	(4.5)	(2.8)	(4.2)	(1.5)	(1.3)	(1.5)	(2.1)	(100.0)				
再導入	2	9	1	8	7	7	2		4	6	3	1	1	2	1	1	55		55	138.65	142.04
(%)	(3.6)	(16.4)	(1.8)	(14.5)	(12.7)	(12.7)	(3.6)		(7.3)	(10.9)	(5.5)	(1.8)	(1.8)	(3.6)	(1.8)	(1.8)	(100.0)				
その他	9	16	15	20	21	23	9	2	6	6	4	13	3	4	4	1	156	1	157	127.48	113.23
(%)	(5.8)	(10.3)	(9.6)	(12.8)	(13.5)	(14.7)	(5.8)	(1.3)	(3.8)	(3.8)	(2.6)	(8.3)	(1.9)	(2.6)	(2.6)	(0.6)	(100.0)				
合計	393	511	499	821	726	589	427	16	305	242	159	197	92	58	67	69	5,171	44	5,215	115.87	120.35
(%)	(7.6)	(9.9)	(9.6)	(15.9)	(14.0)	(11.4)	(8.3)	(0.3)	(5.9)	(4.7)	(3.1)	(3.8)	(1.8)	(1.1)	(1.3)	(1.3)	(100.0)				
記載なし																					
(%)																					
総計	393	511	499	821	726	589	427	16	305	242	159	197	92	58	67	69	5,171	44	5,215	115.87	120.35
(%)	(7.6)	(9.9)	(9.6)	(15.9)	(14.0)	(11.4)	(8.3)	(0.3)	(5.9)	(4.7)	(3.1)	(3.8)	(1.8)	(1.1)	(1.3)	(1.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4273##

(C)Japanese Society for Dialysis Therapy

2016年調査