

表 1215 PTH値(pg/mL) 透析歴(年)別 (腹膜透析患者) PTH測定法:whole-PTH

透析歴	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
2年未満	11	16	16	23	21	26	18		13	20	10	17	6	4	2	3	206	16	222	142.41	123.19
(%)	(5.3)	(7.8)	(7.8)	(11.2)	(10.2)	(12.6)	(8.7)		(6.3)	(9.7)	(4.9)	(8.3)	(2.9)	(1.9)	(1.0)	(1.5)	(100.0)				
2年～	12	22	21	28	24	23	21		14	6	6	4	6		2	4	193	16	209	114.85	114.56
(%)	(6.2)	(11.4)	(10.9)	(14.5)	(12.4)	(11.9)	(10.9)		(7.3)	(3.1)	(3.1)	(2.1)	(3.1)		(1.0)	(2.1)	(100.0)				
5年～	12	8	13	29	16	14	15		8	6	5	4	4	5	2	3	144	5	149	129.35	127.64
(%)	(8.3)	(5.6)	(9.0)	(20.1)	(11.1)	(9.7)	(10.4)		(5.6)	(4.2)	(3.5)	(2.8)	(2.8)	(3.5)	(1.4)	(2.1)	(100.0)				
10年～	2	2	4	7	8	7			3	2	1	3	1		2		42	1	43	123.50	100.95
(%)	(4.8)	(4.8)	(9.5)	(16.7)	(19.0)	(16.7)			(7.1)	(4.8)	(2.4)	(7.1)	(2.4)		(4.8)		(100.0)				
15年～	2		4	5	1	1				1	1	2					17		17	98.00	85.07
(%)	(11.8)		(23.5)	(29.4)	(5.9)	(5.9)				(5.9)	(5.9)	(11.8)					(100.0)				
20年～									1						1		2		2	330.50	235.47
(%)									(50.0)						(50.0)		(100.0)				
25年～							1										1		1	130.00	
(%)							(100.0)										(100.0)				
30年～																					
(%)																					
35年～																					
(%)																					
40年～																					
(%)																					
合計	39	48	58	92	70	71	55		39	35	23	30	17	9	9	10	605	38	643	128.55	120.18
(%)	(6.4)	(7.9)	(9.6)	(15.2)	(11.6)	(11.7)	(9.1)		(6.4)	(5.8)	(3.8)	(5.0)	(2.8)	(1.5)	(1.5)	(1.7)	(100.0)				
不明																					
(%)																					
記載なし																					
(%)																					
総計	39	48	58	92	70	71	55		39	35	23	30	17	9	9	10	605	38	643	128.55	120.18
(%)	(6.4)	(7.9)	(9.6)	(15.2)	(11.6)	(11.7)	(9.1)		(6.4)	(5.8)	(3.8)	(5.0)	(2.8)	(1.5)	(1.5)	(1.7)	(100.0)				
平均	4.54	3.15	4.38	5.00	4.20	3.80	3.78		4.10	3.09	3.52	3.53	3.29	3.56	7.22	3.10	4.05	2.50	3.96		
標準偏差	4.33	2.89	4.40	4.47	3.80	3.88	4.15		4.45	4.08	4.13	4.77	3.22	3.00	7.33	2.60	4.15	2.47	4.08		

数値下のかっこ内は行方向の合計に対する%です。
##D4282##
(C)Japanese Society for Dialysis Therapy
2016年調査