

表 1240 ヘモグロビン濃度(g/dL) 医療機関別 (腹膜透析患者)

医療機関	5未満	5~	6~	7~	8~	9~	10~	11~	12~	13~	14~	合計	記載なし	総計	平均	標準偏差
国公立大学			1	4	18	48	85	97	42	15	1	311	245	556	10.85	1.27
(%)			(0.3)	(1.3)	(5.8)	(15.4)	(27.3)	(31.2)	(13.5)	(4.8)	(0.3)	(100.0)				
私立大学	1		3	11	31	119	219	211	97	28	12	732	417	1,149	10.87	1.34
(%)	(0.1)		(0.4)	(1.5)	(4.2)	(16.3)	(29.9)	(28.8)	(13.3)	(3.8)	(1.6)	(100.0)				
国立病院機構等			5	9	31	41	50	32	14	2	184	43	227	11.01	1.45	
(%)			(2.7)	(4.9)	(16.8)	(22.3)	(27.2)	(17.4)	(7.6)	(1.1)	(100.0)					
県市町村立・国保			18	51	166	326	372	135	48	19	1,135	465	1,600	10.92	1.29	
(%)			(1.6)	(4.5)	(14.6)	(28.7)	(32.8)	(11.9)	(4.2)	(1.7)	(100.0)					
JCHO(旧社会保険)等	2		2	13	27	38	37	28	1	3	151	345	496	10.71	1.48	
(%)	(1.3)		(1.3)	(8.6)	(17.9)	(25.2)	(24.5)	(18.5)	(0.7)	(2.0)	(100.0)					
厚生連等		2	14	18	53	91	112	51	20	6	367	184	551	10.89	1.50	
(%)		(0.5)	(3.8)	(4.9)	(14.4)	(24.8)	(30.5)	(13.9)	(5.4)	(1.6)	(100.0)					
その他公的	1	2	12	43	133	255	269	149	34	13	911	416	1,327	10.94	1.34	
(%)	(0.1)	(0.2)	(1.3)	(4.7)	(14.6)	(28.0)	(29.5)	(16.4)	(3.7)	(1.4)	(100.0)					
私立総合	1	2	2	20	49	87	113	54	19	6	353	40	393	11.01	1.41	
(%)	(0.3)	(0.6)	(0.6)	(5.7)	(13.9)	(24.6)	(32.0)	(15.3)	(5.4)	(1.7)	(100.0)					
私立	1	3	2	24	58	142	313	325	149	45	20	1,082	544	1,626	10.90	1.41
(%)	(0.1)	(0.3)	(0.2)	(2.2)	(5.4)	(13.1)	(28.9)	(30.0)	(13.8)	(4.2)	(1.8)	(100.0)				
私立診療所	1		3	3	22	76	143	207	72	29	7	563	127	690	11.01	1.32
(%)	(0.2)		(0.5)	(0.5)	(3.9)	(13.5)	(25.4)	(36.8)	(12.8)	(5.2)	(1.2)	(100.0)				
総計	3	7	15	95	283	844	1,598	1,793	809	253	89	5,789	2,826	8,615	10.92	1.36
(%)	(0.1)	(0.1)	(0.3)	(1.6)	(4.9)	(14.6)	(27.6)	(31.0)	(14.0)	(4.4)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4344##

(C)Japanese Society for Dialysis Therapy

2016年調査