

表 1241 総コレステロール濃度(mg/dL) 治療方法別 (透析患者全体)

治療方法	100未満	100~	120~	140~	160~	180~	200~	220~	240~	260~	合計	記載なし	総計	平均	標準偏差
施設血液透析	6,910	19,177	34,864	39,904	32,673	20,857	11,045	4,920	2,010	1,470	173,830	59,874	233,704	155.51	36.15
(%)	(4.0)	(11.0)	(20.1)	(23.0)	(18.8)	(12.0)	(6.4)	(2.8)	(1.2)	(0.8)	(100.0)				
血液透析濾過	1,762	5,509	10,814	13,105	11,523	7,664	4,090	1,845	778	460	57,550	17,249	74,799	158.57	35.74
(%)	(3.1)	(9.6)	(18.8)	(22.8)	(20.0)	(13.3)	(7.1)	(3.2)	(1.4)	(0.8)	(100.0)				
血液濾過	2	2	1	9	1	1	2				18	1	19	146.39	32.55
(%)	(11.1)	(11.1)	(5.6)	(50.0)	(5.6)	(5.6)	(11.1)				(100.0)				
血液吸着透析	19	75	198	231	240	151	97	52	18	8	1,089	266	1,355	163.86	35.48
(%)	(1.7)	(6.9)	(18.2)	(21.2)	(22.0)	(13.9)	(8.9)	(4.8)	(1.7)	(0.7)	(100.0)				
在宅血液透析	5	19	32	73	86	83	60	33	18	16	425	192	617	181.82	42.26
(%)	(1.2)	(4.5)	(7.5)	(17.2)	(20.2)	(19.5)	(14.1)	(7.8)	(4.2)	(3.8)	(100.0)				
腹膜透析	60	217	505	815	949	813	526	334	154	155	4,528	4,087	8,615	176.97	41.25
(%)	(1.3)	(4.8)	(11.2)	(18.0)	(21.0)	(18.0)	(11.6)	(7.4)	(3.4)	(3.4)	(100.0)				
合計	8,758	24,999	46,414	54,137	45,472	29,569	15,820	7,184	2,978	2,109	237,440	81,669	319,109	156.74	36.32
(%)	(3.7)	(10.5)	(19.5)	(22.8)	(19.2)	(12.5)	(6.7)	(3.0)	(1.3)	(0.9)	(100.0)				
記載なし															
(%)															
総計	8,758	24,999	46,414	54,137	45,472	29,569	15,820	7,184	2,978	2,109	237,440	81,669	319,109	156.74	36.32
(%)	(3.7)	(10.5)	(19.5)	(22.8)	(19.2)	(12.5)	(6.7)	(3.0)	(1.3)	(0.9)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4410##

(C)Japanese Society for Dialysis Therapy

2016年調査