

表 1243 総コレステロール濃度(mg/dL) 併用の状況別（透析患者全体）

併用の状況	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	8,687	24,762	45,870	53,269	44,495	28,724	15,277	6,836	2,819	1,951	232,690	77,448	310,138	156.35	36.09
(%)	(3.7)	(10.6)	(19.7)	(22.9)	(19.1)	(12.3)	(6.6)	(2.9)	(1.2)	(0.8)	(100.0)				
PD単独	32	140	355	584	705	644	435	282	143	141	3,461	3,470	6,931	180.69	41.82
(%)	(0.9)	(4.0)	(10.3)	(16.9)	(20.4)	(18.6)	(12.6)	(8.1)	(4.1)	(4.1)	(100.0)				
HD(F)単独.PDカテあり	7	15	23	32	18	19	10	7	3	1	135	42	177	160.29	44.24
(%)	(5.2)	(11.1)	(17.0)	(23.7)	(13.3)	(14.1)	(7.4)	(5.2)	(2.2)	(0.7)	(100.0)				
PD.HD併用週1HD(F)	27	65	138	208	216	144	85	41	9	12	945	551	1,496	164.27	36.47
(%)	(2.9)	(6.9)	(14.6)	(22.0)	(22.9)	(15.2)	(9.0)	(4.3)	(1.0)	(1.3)	(100.0)				
PD.HD併用週2HD(F)	2	9	10	27	29	24	7	11	4	1	124	41	165	170.28	38.45
(%)	(1.6)	(7.3)	(8.1)	(21.8)	(23.4)	(19.4)	(5.6)	(8.9)	(3.2)	(0.8)	(100.0)				
PD.HD併用週3HD(F)	2	3	5	8	1	5		3			27	10	37	152.81	39.17
(%)	(7.4)	(11.1)	(18.5)	(29.6)	(3.7)	(18.5)		(11.1)			(100.0)				
PD.HD併用週4HD(F)						1					1		1	183.00	
(%)						(100.0)					(100.0)				
PD.HD併用その他	1	3	8	9	7	7	2	3		1	41	22	63	164.73	41.50
(%)	(2.4)	(7.3)	(19.5)	(22.0)	(17.1)	(17.1)	(4.9)	(7.3)		(2.4)	(100.0)				
合計	8,758	24,997	46,409	54,137	45,471	29,568	15,816	7,183	2,978	2,107	237,424	81,584	319,008	156.74	36.31
(%)	(3.7)	(10.5)	(19.5)	(22.8)	(19.2)	(12.5)	(6.7)	(3.0)	(1.3)	(0.9)	(100.0)				
不明		2	5		1	1	4	1		2	16	6	22	178.56	54.74
(%)		(12.5)	(31.3)		(6.3)	(6.3)	(25.0)	(6.3)		(12.5)	(100.0)				
記載なし												79	79		
(%)															
総計	8,758	24,999	46,414	54,137	45,472	29,569	15,820	7,184	2,978	2,109	237,440	81,669	319,109	156.74	36.32
(%)	(3.7)	(10.5)	(19.5)	(22.8)	(19.2)	(12.5)	(6.7)	(3.0)	(1.3)	(0.9)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4412##

(C)Japanese Society for Dialysis Therapy

2016年調査