

表 1257 総コレステロール濃度(mg/dL) 原疾患別 (血液透析濾過患者 透析回数週3回)

原疾患	100未満	100~	120~	140~	160~	180~	200~	220~	240~	260~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	406	1,372	3,107	4,117	3,973	2,892	1,508	733	318	166	18,592	3,796	22,388	163.26	35.59
(%)	(2.2)	(7.4)	(16.7)	(22.1)	(21.4)	(15.6)	(8.1)	(3.9)	(1.7)	(0.9)	(100.0)				
慢性腎盂腎炎	7	34	94	108	131	82	42	21	12	5	536	91	627	164.33	34.73
(%)	(1.3)	(6.3)	(17.5)	(20.1)	(24.4)	(15.3)	(7.8)	(3.9)	(2.2)	(0.9)	(100.0)				
急速進行性糸球体腎炎	8	19	55	75	100	61	43	22	19	14	416	68	484	173.97	41.94
(%)	(1.9)	(4.6)	(13.2)	(18.0)	(24.0)	(14.7)	(10.3)	(5.3)	(4.6)	(3.4)	(100.0)				
妊娠腎/妊娠中毒症	5	14	52	74	73	68	50	28	9	7	380	79	459	174.12	37.36
(%)	(1.3)	(3.7)	(13.7)	(19.5)	(19.2)	(17.9)	(13.2)	(7.4)	(2.4)	(1.8)	(100.0)				
その他の分類不能の腎炎	5	17	52	58	63	40	30	14	5	1	285	70	355	164.97	35.15
(%)	(1.8)	(6.0)	(18.2)	(20.4)	(22.1)	(14.0)	(10.5)	(4.9)	(1.8)	(0.4)	(100.0)				
多発性囊胞腎	37	176	375	489	462	317	168	68	20	13	2,125	444	2,569	160.85	34.09
(%)	(1.7)	(8.3)	(17.6)	(23.0)	(21.7)	(14.9)	(7.9)	(3.2)	(0.9)	(0.6)	(100.0)				
腎硬化症	128	449	926	1,166	997	625	333	128	51	30	4,833	1,063	5,896	157.80	33.97
(%)	(2.6)	(9.3)	(19.2)	(24.1)	(20.6)	(12.9)	(6.9)	(2.6)	(1.1)	(0.6)	(100.0)				
悪性高血圧	6	33	73	98	102	67	39	19	9	5	451	104	555	165.27	35.11
(%)	(1.3)	(7.3)	(16.2)	(21.7)	(22.6)	(14.9)	(8.6)	(4.2)	(2.0)	(1.1)	(100.0)				
糖尿病性腎症	899	2,563	4,419	4,673	3,616	2,129	1,071	439	193	113	20,115	4,401	24,516	151.69	34.83
(%)	(4.5)	(12.7)	(22.0)	(23.2)	(18.0)	(10.6)	(5.3)	(2.2)	(1.0)	(0.6)	(100.0)				
SLE腎炎	10	31	48	110	88	71	39	32	12	7	448	89	537	168.90	38.02
(%)	(2.2)	(6.9)	(10.7)	(24.6)	(19.6)	(15.8)	(8.7)	(7.1)	(2.7)	(1.6)	(100.0)				
アミロイド腎	1	6	18	9	13	16	11	2	1	1	78	20	98	165.59	37.16
(%)	(1.3)	(7.7)	(23.1)	(11.5)	(16.7)	(20.5)	(14.1)	(2.6)	(1.3)	(1.3)	(100.0)				
痛風腎	3	25	36	38	42	26	11	3	2	4	190	37	227	157.77	35.31
(%)	(1.6)	(13.2)	(18.9)	(20.0)	(22.1)	(13.7)	(5.8)	(1.6)	(1.1)	(2.1)	(100.0)				
先天性代謝異常による腎不全	1	4	8	12	10	11	5	2	1	3	57	16	73	169.11	43.47
(%)	(1.8)	(7.0)	(14.0)	(21.1)	(17.5)	(19.3)	(8.8)	(3.5)	(1.8)	(5.3)	(100.0)				
腎・尿路結核	1	1	2	8	8	2	7		1		30	14	44	169.77	33.79
(%)	(3.3)	(3.3)	(6.7)	(26.7)	(26.7)	(6.7)	(23.3)		(3.3)		(100.0)				
腎・尿路結石	4	8	17	20	18	13	7	5			92	21	113	158.57	34.06
(%)	(4.3)	(8.7)	(18.5)	(21.7)	(19.6)	(14.1)	(7.6)	(5.4)			(100.0)				
腎・尿路腫瘍	6	22	26	36	31	20	9	5	1		156	34	190	153.35	33.75
(%)	(3.8)	(14.1)	(16.7)	(23.1)	(19.9)	(12.8)	(5.8)	(3.2)	(0.6)		(100.0)				
閉塞性尿路障害	1	4	20	26	34	16	19	5			125	26	151	167.21	29.98
(%)	(0.8)	(3.2)	(16.0)	(20.8)	(27.2)	(12.8)	(15.2)	(4.0)			(100.0)				
骨髄腫	1	5	6	4	5	2	4		2		29	8	37	157.62	43.51
(%)	(3.4)	(17.2)	(20.7)	(13.8)	(17.2)	(6.9)	(13.8)		(6.9)		(100.0)				
腎形成不全	3	9	19	30	37	16	14	5	4	4	141	30	171	168.38	38.50
(%)	(2.1)	(6.4)	(13.5)	(21.3)	(26.2)	(11.3)	(9.9)	(3.5)	(2.8)	(2.8)	(100.0)				
不明	131	460	897	1,206	1,064	732	419	190	70	47	5,216	1,315	6,531	160.83	36.02
(%)	(2.5)	(8.8)	(17.2)	(23.1)	(20.4)	(14.0)	(8.0)	(3.6)	(1.3)	(0.9)	(100.0)				
再導入	10	39	90	126	113	63	44	29	9	10	533	100	633	165.01	38.16
(%)	(1.9)	(7.3)	(16.9)	(23.6)	(21.2)	(11.8)	(8.3)	(5.4)	(1.7)	(1.9)	(100.0)				
その他	41	112	257	300	276	202	118	44	16	14	1,380	340	1,720	160.83	36.29
(%)	(3.0)	(8.1)	(18.6)	(21.7)	(20.0)	(14.6)	(8.6)	(3.2)	(1.2)	(1.0)	(100.0)				
合計	1,714	5,403	10,597	12,783	11,256	7,471	3,991	1,794	755	444	56,208	12,166	68,374	158.50	35.72
(%)	(3.0)	(9.6)	(18.9)	(22.7)	(20.0)	(13.3)	(7.1)	(3.2)	(1.3)	(0.8)	(100.0)				
記載なし															
(%)															
総計	1,714	5,403	10,597	12,783	11,256	7,471	3,991	1,794	755	444	56,208	12,166	68,374	158.50	35.72
(%)	(3.0)	(9.6)	(18.9)	(22.7)	(20.0)	(13.3)	(7.1)	(3.2)	(1.3)	(0.8)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4433##

(C)Japanese Society for Dialysis Therapy

2016年調査