

表 1265 HDL-C濃度(mg/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	20未満	20~	30~	40~	50~	60~	70~	80~	90~	100~	合計	記載なし	総計	平均	標準偏差
施設血液透析	644	13,327	39,366	43,706	31,574	18,067	8,806	3,897	1,773	1,530	162,690	71,014	233,704	48.28	16.32
(%)	(0.4)	(8.2)	(24.2)	(26.9)	(19.4)	(11.1)	(5.4)	(2.4)	(1.1)	(0.9)	(100.0)				
Off-line HDF	13	256	691	794	650	373	197	86	31	37	3,128	1,509	4,637	49.40	16.85
(%)	(0.4)	(8.2)	(22.1)	(25.4)	(20.8)	(11.9)	(6.3)	(2.7)	(1.0)	(1.2)	(100.0)				
On-line HDF	134	2,779	9,357	11,456	9,011	5,781	3,095	1,546	652	471	44,282	14,834	59,116	50.65	16.87
(%)	(0.3)	(6.3)	(21.1)	(25.9)	(20.3)	(13.1)	(7.0)	(3.5)	(1.5)	(1.1)	(100.0)				
Push/Pull HDF		1	5	13	10	9	2	1	1		42	24	66	53.19	14.03
(%)		(2.4)	(11.9)	(31.0)	(23.8)	(21.4)	(4.8)	(2.4)	(2.4)		(100.0)				
バイオフィルトレーション		12	49	61	29	19	14	2	3	1	190	62	252	48.17	15.87
(%)		(6.3)	(25.8)	(32.1)	(15.3)	(10.0)	(7.4)	(1.1)	(1.6)	(0.5)	(100.0)				
IHDF	17	654	1,916	2,185	1,618	876	430	188	82	55	8,021	2,707	10,728	48.19	15.65
(%)	(0.2)	(8.2)	(23.9)	(27.2)	(20.2)	(10.9)	(5.4)	(2.3)	(1.0)	(0.7)	(100.0)				
血液濾過		4	7	3	1	2					17	2	19	37.00	12.49
(%)		(23.5)	(41.2)	(17.6)	(5.9)	(11.8)					(100.0)				
血液吸着透析 (リクセル等使用)	3	47	187	256	237	177	76	27	14	11	1,035	320	1,355	52.21	16.62
(%)	(0.3)	(4.5)	(18.1)	(24.7)	(22.9)	(17.1)	(7.3)	(2.6)	(1.4)	(1.1)	(100.0)				
在宅血液透析	3	16	58	92	90	70	30	24	13	16	412	205	617	56.80	20.59
(%)	(0.7)	(3.9)	(14.1)	(22.3)	(21.8)	(17.0)	(7.3)	(5.8)	(3.2)	(3.9)	(100.0)				
腹膜透析 (CAPD)	7	147	512	595	464	309	131	88	32	27	2,312	2,661	4,973	50.40	17.87
(%)	(0.3)	(6.4)	(22.1)	(25.7)	(20.1)	(13.4)	(5.7)	(3.8)	(1.4)	(1.2)	(100.0)				
腹膜透析 (APD)	8	93	269	306	213	157	82	42	29	25	1,224	1,413	2,637	50.87	18.78
(%)	(0.7)	(7.6)	(22.0)	(25.0)	(17.4)	(12.8)	(6.7)	(3.4)	(2.4)	(2.0)	(100.0)				
腹膜透析 (CCPD)		49	140	128	107	55	31	19	6	4	539	466	1,005	48.67	17.65
(%)		(9.1)	(26.0)	(23.7)	(19.9)	(10.2)	(5.8)	(3.5)	(1.1)	(0.7)	(100.0)				
合計	829	17,385	52,557	59,595	44,004	25,895	12,894	5,920	2,636	2,177	223,892	95,217	319,109	48.83	16.49
(%)	(0.4)	(7.8)	(23.5)	(26.6)	(19.7)	(11.6)	(5.8)	(2.6)	(1.2)	(1.0)	(100.0)				
記載なし															
(%)															
総計	829	17,385	52,557	59,595	44,004	25,895	12,894	5,920	2,636	2,177	223,892	95,217	319,109	48.83	16.49
(%)	(0.4)	(7.8)	(23.5)	(26.6)	(19.7)	(11.6)	(5.8)	(2.6)	(1.2)	(1.0)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

##D4511##

(C)Japanese Society for Dialysis Therapy

2016年調査