

表 1275 HDL-C濃度(mg/dL) 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	20未満	20~	30~	40~	50~	60~	70~	80~	90~	100~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	121	2,370	8,659	11,548	9,189	5,883	3,052	1,382	619	475	43,298	13,874	57,172	51.04	16.60
(%)	(0.3)	(5.5)	(20.0)	(26.7)	(21.2)	(13.6)	(7.0)	(3.2)	(1.4)	(1.1)	(100.0)				
慢性腎盂腎炎	6	69	222	350	324	191	93	52	17	19	1,343	525	1,868	52.15	16.89
(%)	(0.4)	(5.1)	(16.5)	(26.1)	(24.1)	(14.2)	(6.9)	(3.9)	(1.3)	(1.4)	(100.0)				
急速進行性糸球体腎炎	2	39	201	275	298	209	123	59	31	33	1,270	417	1,687	55.67	18.20
(%)	(0.2)	(3.1)	(15.8)	(21.7)	(23.5)	(16.5)	(9.7)	(4.6)	(2.4)	(2.6)	(100.0)				
妊娠腎/妊娠中毒症		14	102	140	161	110	59	42	15	12	655	194	849	55.98	17.51
(%)		(2.1)	(15.6)	(21.4)	(24.6)	(16.8)	(9.0)	(6.4)	(2.3)	(1.8)	(100.0)				
その他の分類不能の腎炎	1	25	142	177	169	105	42	24	9	6	700	202	902	51.46	15.42
(%)	(0.1)	(3.6)	(20.3)	(25.3)	(24.1)	(15.0)	(6.0)	(3.4)	(1.3)	(0.9)	(100.0)				
多発性囊胞腎	20	297	1,178	1,566	1,290	783	361	133	66	37	5,731	1,817	7,548	50.19	15.51
(%)	(0.3)	(5.2)	(20.6)	(27.3)	(22.5)	(13.7)	(6.3)	(2.3)	(1.2)	(0.6)	(100.0)				
腎硬化症	61	1,147	3,715	4,562	3,263	1,797	880	355	177	134	16,091	5,427	21,518	48.59	15.89
(%)	(0.4)	(7.1)	(23.1)	(28.4)	(20.3)	(11.2)	(5.5)	(2.2)	(1.1)	(0.8)	(100.0)				
悪性高血圧	5	74	318	384	268	151	80	31	16	20	1,347	430	1,777	49.58	16.77
(%)	(0.4)	(5.5)	(23.6)	(28.5)	(19.9)	(11.2)	(5.9)	(2.3)	(1.2)	(1.5)	(100.0)				
糖尿病性腎症	315	7,337	18,325	16,992	10,600	5,244	2,289	995	410	438	62,945	21,523	84,468	45.01	15.30
(%)	(0.5)	(11.7)	(29.1)	(27.0)	(16.8)	(8.3)	(3.6)	(1.6)	(0.7)	(0.7)	(100.0)				
SLE腎炎	3	32	150	232	235	173	109	62	34	14	1,044	326	1,370	55.88	17.70
(%)	(0.3)	(3.1)	(14.4)	(22.2)	(22.5)	(16.6)	(10.4)	(5.9)	(3.3)	(1.3)	(100.0)				
アミロイド腎	1	7	35	43	49	35	18	7	13	4	212	68	280	56.20	19.67
(%)	(0.5)	(3.3)	(16.5)	(20.3)	(23.1)	(16.5)	(8.5)	(3.3)	(6.1)	(1.9)	(100.0)				
痛風腎	1	46	109	143	108	62	28	19	8	8	532	168	700	49.58	17.39
(%)	(0.2)	(8.6)	(20.5)	(26.9)	(20.3)	(11.7)	(5.3)	(3.6)	(1.5)	(1.5)	(100.0)				
先天性代謝異常による腎不全	1	4	21	36	27	17	6	3	3	5	123	46	169	53.55	20.36
(%)	(0.8)	(3.3)	(17.1)	(29.3)	(22.0)	(13.8)	(4.9)	(2.4)	(2.4)	(4.1)	(100.0)				
腎・尿路結核		7	14	21	20	12	3	2	2	2	81	25	106	49.89	15.42
(%)		(8.6)	(17.3)	(25.9)	(24.7)	(14.8)	(3.7)	(2.5)	(2.5)	(2.5)	(100.0)				
腎・尿路結石		17	57	79	62	30	21	5	3	6	280	104	384	50.30	17.40
(%)		(6.1)	(20.4)	(28.2)	(22.1)	(10.7)	(7.5)	(1.8)	(1.1)	(2.1)	(100.0)				
腎・尿路腫瘍	3	40	117	127	109	51	26	12	5	6	496	148	644	48.68	16.20
(%)	(0.6)	(8.1)	(23.6)	(25.6)	(22.0)	(10.3)	(5.2)	(2.4)	(1.0)	(1.2)	(100.0)				
閉塞性尿路障害		11	69	92	90	46	22	10	6	10	356	109	465	53.14	19.10
(%)		(3.1)	(19.4)	(25.8)	(25.3)	(12.9)	(6.2)	(2.8)	(1.7)	(2.8)	(100.0)				
骨髄腫		12	26	38	33	20	13	5	1	2	150	63	213	50.94	16.30
(%)		(8.0)	(17.3)	(25.3)	(22.0)	(13.3)	(8.7)	(3.3)	(0.7)	(1.3)	(100.0)				
腎形成不全		15	55	85	59	35	34	6	2	4	295	73	368	51.54	16.75
(%)		(5.1)	(18.6)	(28.8)	(20.0)	(11.9)	(11.5)	(2.0)	(0.7)	(1.4)	(100.0)				
不明	57	968	3,340	3,899	3,045	1,776	877	390	192	158	14,702	5,776	20,478	49.51	16.50
(%)	(0.4)	(6.6)	(22.7)	(26.5)	(20.7)	(12.1)	(6.0)	(2.7)	(1.3)	(1.1)	(100.0)				
再導入	2	46	201	230	187	110	63	27	18	8	892	272	1,164	50.68	16.68
(%)	(0.2)	(5.2)	(22.5)	(25.8)	(21.0)	(12.3)	(7.1)	(3.0)	(2.0)	(0.9)	(100.0)				
その他	17	273	845	998	849	503	252	132	58	52	3,979	1,427	5,406	50.43	17.31
(%)	(0.4)	(6.9)	(21.2)	(25.1)	(21.3)	(12.6)	(6.3)	(3.3)	(1.5)	(1.3)	(100.0)				
合計	616	12,850	37,901	42,017	30,435	17,343	8,451	3,753	1,705	1,451	156,522	53,014	209,536	48.26	16.29
(%)	(0.4)	(8.2)	(24.2)	(26.8)	(19.4)	(11.1)	(5.4)	(2.4)	(1.1)	(0.9)	(100.0)				
記載なし															
(%)															
総計	616	12,850	37,901	42,017	30,435	17,343	8,451	3,753	1,705	1,451	156,522	53,014	209,536	48.26	16.29
(%)	(0.4)	(8.2)	(24.2)	(26.8)	(19.4)	(11.1)	(5.4)	(2.4)	(1.1)	(0.9)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4523##

(C)Japanese Society for Dialysis Therapy

2016年調査