

表 1288 Non-HDL-C濃度(mg/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	50未満	50~	70~	90~	110~	130~	150~	170~	190~	210~	合計	記載なし	総計	平均	標準偏差
施設血液透析	2,660	13,513	29,825	35,280	28,272	17,331	8,873	3,908	1,515	1,033	142,210	91,494	233,704	107.19	33.70
(%)	(1.9)	(9.5)	(21.0)	(24.8)	(19.9)	(12.2)	(6.2)	(2.7)	(1.1)	(0.7)	(100.0)				
Off-line HDF	54	271	584	671	574	311	183	58	28	22	2,756	1,881	4,637	106.53	33.60
(%)	(2.0)	(9.8)	(21.2)	(24.3)	(20.8)	(11.3)	(6.6)	(2.1)	(1.0)	(0.8)	(100.0)				
On-line HDF	664	3,290	7,762	9,633	8,058	5,043	2,588	1,060	447	214	38,759	20,357	59,116	108.49	33.12
(%)	(1.7)	(8.5)	(20.0)	(24.9)	(20.8)	(13.0)	(6.7)	(2.7)	(1.2)	(0.6)	(100.0)				
Push/Pull HDF	1	3	6	10	8	7	5	2			42	24	66	114.38	32.84
(%)	(2.4)	(7.1)	(14.3)	(23.8)	(19.0)	(16.7)	(11.9)	(4.8)			(100.0)				
バイオフィルトレーション	5	17	39	32	35	13	12	8	2	4	167	85	252	108.66	37.93
(%)	(3.0)	(10.2)	(23.4)	(19.2)	(21.0)	(7.8)	(7.2)	(4.8)	(1.2)	(2.4)	(100.0)				
IHDF	112	665	1,519	1,662	1,369	827	413	182	64	44	6,857	3,871	10,728	106.58	32.98
(%)	(1.6)	(9.7)	(22.2)	(24.2)	(20.0)	(12.1)	(6.0)	(2.7)	(0.9)	(0.6)	(100.0)				
血液濾過		3	1	2	6	3	2				17	2	19	111.59	30.75
(%)		(17.6)	(5.9)	(11.8)	(35.3)	(17.6)	(11.8)				(100.0)				
血液吸着透析 (リクセル等使用)	13	72	179	213	214	125	76	34	11	8	945	410	1,355	111.29	33.40
(%)	(1.4)	(7.6)	(18.9)	(22.5)	(22.6)	(13.2)	(8.0)	(3.6)	(1.2)	(0.8)	(100.0)				
在宅血液透析	5	15	34	88	77	73	44	17	7	7	367	250	617	122.07	35.49
(%)	(1.4)	(4.1)	(9.3)	(24.0)	(21.0)	(19.9)	(12.0)	(4.6)	(1.9)	(1.9)	(100.0)				
腹膜透析 (CAPD)	10	56	202	378	420	357	229	151	58	62	1,923	3,050	4,973	128.69	39.65
(%)	(0.5)	(2.9)	(10.5)	(19.7)	(21.8)	(18.6)	(11.9)	(7.9)	(3.0)	(3.2)	(100.0)				
腹膜透析 (APD)	12	45	132	239	250	178	129	72	20	17	1,094	1,543	2,637	121.88	36.19
(%)	(1.1)	(4.1)	(12.1)	(21.8)	(22.9)	(16.3)	(11.8)	(6.6)	(1.8)	(1.6)	(100.0)				
腹膜透析 (CCPD)	2	21	56	90	106	90	49	32	11	13	470	535	1,005	125.18	38.34
(%)	(0.4)	(4.5)	(11.9)	(19.1)	(22.6)	(19.1)	(10.4)	(6.8)	(2.3)	(2.8)	(100.0)				
合計	3,538	17,971	40,339	48,298	39,389	24,358	12,603	5,524	2,163	1,424	195,607	123,502	319,109	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				
記載なし (%)															
総計	3,538	17,971	40,339	48,298	39,389	24,358	12,603	5,524	2,163	1,424	195,607	123,502	319,109	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

##D4611##

(C)Japanese Society for Dialysis Therapy

2016年調査