

表 1289 Non-HDL-C濃度(mg/dL) 併用の状況別 (透析患者全体)

併用の状況	50未満	50~	70~	90~	110~	130~	150~	170~	190~	210~	合計	記載なし	総計	平均	標準偏差
HD(F)単独	3,508	17,835	39,922	47,551	38,582	23,709	12,182	5,256	2,071	1,330	191,946	118,192	310,138	107.47	33.57
(%)	(1.8)	(9.3)	(20.8)	(24.8)	(20.1)	(12.4)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				
PD単独	15	81	276	527	587	486	327	206	75	82	2,662	4,269	6,931	128.20	39.08
(%)	(0.6)	(3.0)	(10.4)	(19.8)	(22.1)	(18.3)	(12.3)	(7.7)	(2.8)	(3.1)	(100.0)				
HD(F)単独.PDカテあり	4	12	20	25	18	16	9	7	1	1	113	64	177	110.14	37.15
(%)	(3.5)	(10.6)	(17.7)	(22.1)	(15.9)	(14.2)	(8.0)	(6.2)	(0.9)	(0.9)	(100.0)				
PD.HD併用週1 HD(F)	8	37	101	163	167	130	68	45	14	8	741	755	1,496	119.29	35.59
(%)	(1.1)	(5.0)	(13.6)	(22.0)	(22.5)	(17.5)	(9.2)	(6.1)	(1.9)	(1.1)	(100.0)				
PD.HD併用週2 HD(F)	2	2	11	15	23	9	10	6	2	1	81	84	165	120.95	38.14
(%)	(2.5)	(2.5)	(13.6)	(18.5)	(28.4)	(11.1)	(12.3)	(7.4)	(2.5)	(1.2)	(100.0)				
PD.HD併用週3 HD(F)	1	2	4	5	2	3	2	1			20	17	37	107.40	38.12
(%)	(5.0)	(10.0)	(20.0)	(25.0)	(10.0)	(15.0)	(10.0)	(5.0)			(100.0)				
PD.HD併用週4 HD(F)											1	1			
(%)											(100.0)				
PD.HD併用その他		2	5	11	9	5	4	2		1	39	24	63	119.10	38.11
(%)		(5.1)	(12.8)	(28.2)	(23.1)	(12.8)	(10.3)	(5.1)		(2.6)	(100.0)				
合計	3,538	17,971	40,339	48,297	39,388	24,358	12,602	5,523	2,163	1,423	195,602	123,406	319,008	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				
不明				1	1		1	1		1	5	17	22	152.40	49.07
(%)				(20.0)	(20.0)		(20.0)	(20.0)		(20.0)	(100.0)				
記載なし											79	79			
(%)											(100.0)				
総計	3,538	17,971	40,339	48,298	39,389	24,358	12,603	5,524	2,163	1,424	195,607	123,502	319,109	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4612##

(C)Japanese Society for Dialysis Therapy
2016年調査