

表 1291 Non-HDL-C濃度(mg/dL) 年齢(歳)別 (透析患者全体)

年齢	50未満	50 ~	70 ~	90 ~	110 ~	130 ~	150 ~	170 ~	190 ~	210 ~	合計	記載なし	総計	平均	標準偏差
15歳未満		1	2	1	3	2	4	1		1	15	78	93	130.60	43.07
(%)		(6.7)	(13.3)	(6.7)	(20.0)	(13.3)	(26.7)	(6.7)		(6.7)	(100.0)				
15歳 ~	11	55	112	126	101	53	30	9	7	4	508	403	911	105.23	35.21
(%)	(2.2)	(10.8)	(22.0)	(24.8)	(19.9)	(10.4)	(5.9)	(1.8)	(1.4)	(0.8)	(100.0)				
30歳 ~	114	637	1,520	1,974	1,800	1,140	653	264	131	91	8,324	5,098	13,422	112.06	34.76
(%)	(1.4)	(7.7)	(18.3)	(23.7)	(21.6)	(13.7)	(7.8)	(3.2)	(1.6)	(1.1)	(100.0)				
45歳 ~	559	2,782	6,350	8,330	7,324	5,133	2,902	1,305	540	370	35,595	21,749	57,344	112.48	35.30
(%)	(1.6)	(7.8)	(17.8)	(23.4)	(20.6)	(14.4)	(8.2)	(3.7)	(1.5)	(1.0)	(100.0)				
60歳 ~	1,588	7,930	17,741	21,316	17,721	11,032	5,775	2,560	967	639	87,269	54,360	141,629	108.20	33.90
(%)	(1.8)	(9.1)	(20.3)	(24.4)	(20.3)	(12.6)	(6.6)	(2.9)	(1.1)	(0.7)	(100.0)				
75歳 ~	1,197	6,193	13,737	15,559	11,768	6,626	3,108	1,323	487	309	60,307	39,306	99,613	104.26	32.18
(%)	(2.0)	(10.3)	(22.8)	(25.8)	(19.5)	(11.0)	(5.2)	(2.2)	(0.8)	(0.5)	(100.0)				
90歳 ~	69	373	877	992	672	372	131	62	31	10	3,589	2,506	6,095	101.93	30.39
(%)	(1.9)	(10.4)	(24.4)	(27.6)	(18.7)	(10.4)	(3.7)	(1.7)	(0.9)	(0.3)	(100.0)				
合計	3,538	17,971	40,339	48,298	39,389	24,358	12,603	5,524	2,163	1,424	195,607	123,500	319,107	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				
不明															
(%)															
記載なし													2	2	
(%)															
総計	3,538	17,971	40,339	48,298	39,389	24,358	12,603	5,524	2,163	1,424	195,607	123,502	319,109	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				
平均	69.16	69.27	69.10	68.52	67.71	66.87	65.83	65.87	65.04	64.43	68.04	68.32	68.15		
標準偏差	11.97	12.20	12.31	12.40	12.43	12.39	12.38	12.23	12.74	12.84	12.41	12.59	12.48		

数値下のカッコ内は行方向の合計に対する%です。

##D4614##

(C)Japanese Society for Dialysis Therapy

2016年調査