

表 1293 Non-HDL-C濃度(mg/dL) 原疾患別 (透析患者全体)

原疾患	50未満	50~	70~	90~	110~	130~	150~	170~	190~	210~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	891	4,606	11,085	14,292	12,017	7,678	3,912	1,680	652	393	57,206	34,577	91,783	109.50	33.27
(%)	(1.6)	(8.1)	(19.4)	(25.0)	(21.0)	(13.4)	(6.8)	(2.9)	(1.1)	(0.7)	(100.0)				
慢性腎盂腎炎	29	125	332	435	365	247	128	63	20	12	1,756	1,140	2,896	110.75	33.40
(%)	(1.7)	(7.1)	(18.9)	(24.8)	(20.8)	(14.1)	(7.3)	(3.6)	(1.1)	(0.7)	(100.0)				
急速進行性糸球体腎炎	14	83	239	392	322	247	132	79	27	33	1,568	970	2,538	118.16	37.34
(%)	(0.9)	(5.3)	(15.2)	(25.0)	(20.5)	(15.8)	(8.4)	(5.0)	(1.7)	(2.1)	(100.0)				
妊娠腎/妊娠中毒症	6	51	150	219	203	160	95	50	11	14	959	524	1,483	118.73	34.79
(%)	(0.6)	(5.3)	(15.6)	(22.8)	(21.2)	(16.7)	(9.9)	(5.2)	(1.1)	(1.5)	(100.0)				
その他の分類不能の腎炎	13	77	170	233	200	124	65	32	7	2	923	531	1,454	109.21	32.36
(%)	(1.4)	(8.3)	(18.4)	(25.2)	(21.7)	(13.4)	(7.0)	(3.5)	(0.8)	(0.2)	(100.0)				
多発性囊胞腎	74	485	1,291	1,834	1,689	996	542	197	57	42	7,207	4,264	11,471	111.05	31.78
(%)	(1.0)	(6.7)	(17.9)	(25.4)	(23.4)	(13.8)	(7.5)	(2.7)	(0.8)	(0.6)	(100.0)				
腎硬化症	294	1,645	4,033	5,034	4,018	2,456	1,138	545	201	116	19,480	12,170	31,650	107.81	32.60
(%)	(1.5)	(8.4)	(20.7)	(25.8)	(20.6)	(12.6)	(5.8)	(2.8)	(1.0)	(0.6)	(100.0)				
悪性高血圧	19	130	294	413	356	221	129	62	29	18	1,671	1,050	2,721	112.38	34.59
(%)	(1.1)	(7.8)	(17.6)	(24.7)	(21.3)	(13.2)	(7.7)	(3.7)	(1.7)	(1.1)	(100.0)				
糖尿病性腎症	1,676	8,270	16,920	18,378	14,165	8,505	4,454	1,924	833	563	75,688	47,969	123,657	105.22	34.15
(%)	(2.2)	(10.9)	(22.4)	(24.3)	(18.7)	(11.2)	(5.9)	(2.5)	(1.1)	(0.7)	(100.0)				
SLE腎炎	18	102	243	326	317	186	135	58	17	12	1,414	808	2,222	113.57	34.37
(%)	(1.3)	(7.2)	(17.2)	(23.1)	(22.4)	(13.2)	(9.5)	(4.1)	(1.2)	(0.8)	(100.0)				
アミロイド腎	5	26	61	56	49	30	19	9	2	2	259	179	438	106.71	35.43
(%)	(1.9)	(10.0)	(23.6)	(21.6)	(18.9)	(11.6)	(7.3)	(3.5)	(0.8)	(0.8)	(100.0)				
痛風腎	12	70	135	179	134	79	43	12	6	8	678	378	1,056	106.23	33.44
(%)	(1.8)	(10.3)	(19.9)	(26.4)	(19.8)	(11.7)	(6.3)	(1.8)	(0.9)	(1.2)	(100.0)				
先天性代謝異常による腎不全	4	17	32	37	38	18	11	5	1	3	166	115	281	107.91	36.71
(%)	(2.4)	(10.2)	(19.3)	(22.3)	(22.9)	(10.8)	(6.6)	(3.0)	(0.6)	(1.8)	(100.0)				
腎・尿路結核	2	7	25	24	22	5	9	6	1		101	67	168	108.01	34.19
(%)	(2.0)	(6.9)	(24.8)	(23.8)	(21.8)	(5.0)	(8.9)	(5.9)	(1.0)		(100.0)				
腎・尿路結石	5	39	63	82	73	45	23	6	4	2	342	236	578	106.75	32.91
(%)	(1.5)	(11.4)	(18.4)	(24.0)	(21.3)	(13.2)	(6.7)	(1.8)	(1.2)	(0.6)	(100.0)				
腎・尿路腫瘍	12	53	121	159	146	55	34	14	5	2	601	358	959	105.56	31.11
(%)	(2.0)	(8.8)	(20.1)	(26.5)	(24.3)	(9.2)	(5.7)	(2.3)	(0.8)	(0.3)	(100.0)				
閉塞性尿路障害	4	32	91	120	117	64	35	8	8	3	482	261	743	110.81	32.05
(%)	(0.8)	(6.6)	(18.9)	(24.9)	(24.3)	(13.3)	(7.3)	(1.7)	(1.7)	(0.6)	(100.0)				
骨髄腫	5	15	30	39	35	27	15	10	3	4	183	140	323	115.38	41.58
(%)	(2.7)	(8.2)	(16.4)	(21.3)	(19.1)	(14.8)	(8.2)	(5.5)	(1.6)	(2.2)	(100.0)				
腎形成不全	6	36	66	93	105	47	31	14	4	5	407	257	664	112.02	35.11
(%)	(1.5)	(8.8)	(16.2)	(22.9)	(25.8)	(11.5)	(7.6)	(3.4)	(1.0)	(1.2)	(100.0)				
不明	323	1,569	3,696	4,474	3,739	2,304	1,189	545	203	126	18,168	13,244	31,412	108.44	33.77
(%)	(1.8)	(8.6)	(20.3)	(24.6)	(20.6)	(12.7)	(6.5)	(3.0)	(1.1)	(0.7)	(100.0)				
再導入	21	105	269	316	276	188	121	48	15	16	1,375	799	2,174	112.17	35.97
(%)	(1.5)	(7.6)	(19.6)	(23.0)	(20.1)	(13.7)	(8.8)	(3.5)	(1.1)	(1.2)	(100.0)				
その他	105	428	993	1,163	1,003	676	343	157	57	48	4,973	3,465	8,438	109.24	34.89
(%)	(2.1)	(8.6)	(20.0)	(23.4)	(20.2)	(13.6)	(6.9)	(3.2)	(1.1)	(1.0)	(100.0)				
合計	3,538	17,971	40,339	48,298	39,389	24,358	12,603	5,524	2,163	1,424	195,607	123,502	319,109	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				
記載なし															
(%)															
総計	3,538	17,971	40,339	48,298	39,389	24,358	12,603	5,524	2,163	1,424	195,607	123,502	319,109	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4616##

(C)Japanese Society for Dialysis Therapy

2016年調査