

表 1294 Non-HDL-C濃度(mg/dL) 医療機関別 (透析患者全体)

医療機関	50未満	50~	70~	90~	110~	130~	150~	170~	190~	210~	合計	記載なし	総計	平均	標準偏差
国公立大学	9	22	55	75	60	42	28	19	9	4	323	801	1,124	114.60	38.76
(%)	(2.8)	(6.8)	(17.0)	(23.2)	(18.6)	(13.0)	(8.7)	(5.9)	(2.8)	(1.2)	(100.0)				
私立大学	26	108	251	321	293	185	119	48	27	21	1,399	2,287	3,686	113.09	36.75
(%)	(1.9)	(7.7)	(17.9)	(22.9)	(20.9)	(13.2)	(8.5)	(3.4)	(1.9)	(1.5)	(100.0)				
国立病院機構等	10	26	60	80	62	48	24	13	8	6	337	574	911	112.00	38.76
(%)	(3.0)	(7.7)	(17.8)	(23.7)	(18.4)	(14.2)	(7.1)	(3.9)	(2.4)	(1.8)	(100.0)				
県市町村立・国保	239	1,192	2,344	2,546	1,994	1,093	589	261	103	97	10,458	9,290	19,748	104.71	34.69
(%)	(2.3)	(11.4)	(22.4)	(24.3)	(19.1)	(10.5)	(5.6)	(2.5)	(1.0)	(0.9)	(100.0)				
JCHO(旧社会保険)等	32	208	428	498	432	295	140	86	23	23	2,165	1,600	3,765	109.60	35.35
(%)	(1.5)	(9.6)	(19.8)	(23.0)	(20.0)	(13.6)	(6.5)	(4.0)	(1.1)	(1.1)	(100.0)				
厚生連等	106	526	1,249	1,429	1,200	794	358	184	74	57	5,977	3,408	9,385	108.50	34.29
(%)	(1.8)	(8.8)	(20.9)	(23.9)	(20.1)	(13.3)	(6.0)	(3.1)	(1.2)	(1.0)	(100.0)				
その他公的	105	518	1,172	1,369	1,128	681	357	172	61	40	5,603	5,255	10,858	107.62	34.18
(%)	(1.9)	(9.2)	(20.9)	(24.4)	(20.1)	(12.2)	(6.4)	(3.1)	(1.1)	(0.7)	(100.0)				
私立総合	110	474	1,047	1,222	921	565	293	126	39	36	4,833	2,980	7,813	105.76	33.64
(%)	(2.3)	(9.8)	(21.7)	(25.3)	(19.1)	(11.7)	(6.1)	(2.6)	(0.8)	(0.7)	(100.0)				
私立	1,353	5,793	11,929	13,776	10,906	6,829	3,425	1,532	618	377	56,538	35,051	91,589	106.20	34.14
(%)	(2.4)	(10.2)	(21.1)	(24.4)	(19.3)	(12.1)	(6.1)	(2.7)	(1.1)	(0.7)	(100.0)				
私立診療所	1,548	9,104	21,804	26,982	22,393	13,826	7,270	3,083	1,201	763	107,974	62,256	170,230	108.87	33.25
(%)	(1.4)	(8.4)	(20.2)	(25.0)	(20.7)	(12.8)	(6.7)	(2.9)	(1.1)	(0.7)	(100.0)				
総計	3,538	17,971	40,339	48,298	39,389	24,358	12,603	5,524	2,163	1,424	195,607	123,502	319,109	107.80	33.76
(%)	(1.8)	(9.2)	(20.6)	(24.7)	(20.1)	(12.5)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4617##

(C)Japanese Society for Dialysis Therapy

2016年調査