

表 1298 Non-HDL-C濃度(mg/dL) 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	50未満	50~	70~	90~	110~	130~	150~	170~	190~	210~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	617	3,162	7,508	9,483	7,816	4,956	2,454	1,084	399	249	37,728	19,444	57,172	108.66	33.13
(%)	(1.6)	(8.4)	(19.9)	(25.1)	(20.7)	(13.1)	(6.5)	(2.9)	(1.1)	(0.7)	(100.0)				
慢性腎盂腎炎	22	91	224	276	249	176	85	39	14	7	1,183	685	1,868	110.50	33.64
(%)	(1.9)	(7.7)	(18.9)	(23.3)	(21.0)	(14.9)	(7.2)	(3.3)	(1.2)	(0.6)	(100.0)				
急速進行性糸球体腎炎	10	57	180	269	226	176	97	55	20	20	1,110	577	1,687	117.79	36.61
(%)	(0.9)	(5.1)	(16.2)	(24.2)	(20.4)	(15.9)	(8.7)	(5.0)	(1.8)	(1.8)	(100.0)				
妊娠腎/妊娠中毒症	4	37	89	121	130	91	49	32	8	7	568	281	849	117.86	34.72
(%)	(0.7)	(6.5)	(15.7)	(21.3)	(22.9)	(16.0)	(8.6)	(5.6)	(1.4)	(1.2)	(100.0)				
その他の分類不能の腎炎	10	56	107	160	132	80	40	23	5	1	614	288	902	108.77	32.75
(%)	(1.6)	(9.1)	(17.4)	(26.1)	(21.5)	(13.0)	(6.5)	(3.7)	(0.8)	(0.2)	(100.0)				
多発性囊胞腎	51	331	886	1,293	1,178	693	386	141	38	30	5,027	2,521	7,548	111.26	31.68
(%)	(1.0)	(6.6)	(17.6)	(25.7)	(23.4)	(13.8)	(7.7)	(2.8)	(0.8)	(0.6)	(100.0)				
腎硬化症	213	1,268	2,974	3,693	2,891	1,728	819	369	131	72	14,158	7,360	21,518	106.97	32.17
(%)	(1.5)	(9.0)	(21.0)	(26.1)	(20.4)	(12.2)	(5.8)	(2.6)	(0.9)	(0.5)	(100.0)				
悪性高血圧	15	101	208	290	245	154	90	40	16	11	1,170	607	1,777	111.14	34.17
(%)	(1.3)	(8.6)	(17.8)	(24.8)	(20.9)	(13.2)	(7.7)	(3.4)	(1.4)	(0.9)	(100.0)				
糖尿病性腎症	1,261	6,174	12,445	13,442	10,205	6,122	3,186	1,357	598	421	55,211	29,257	84,468	104.81	34.17
(%)	(2.3)	(11.2)	(22.5)	(24.3)	(18.5)	(11.1)	(5.8)	(2.5)	(1.1)	(0.8)	(100.0)				
SLE腎炎	10	62	164	224	205	127	82	39	11	10	934	436	1,370	113.69	34.35
(%)	(1.1)	(6.6)	(17.6)	(24.0)	(21.9)	(13.6)	(8.8)	(4.2)	(1.2)	(1.1)	(100.0)				
アミロイド腎	4	21	44	39	36	21	12	3	2	2	182	98	280	103.85	34.33
(%)	(2.2)	(11.5)	(24.2)	(21.4)	(19.8)	(11.5)	(6.6)	(1.6)	(1.1)	(1.1)	(100.0)				
痛風腎	10	48	98	119	90	55	26	7	4	6	463	237	700	104.96	33.27
(%)	(2.2)	(10.4)	(21.2)	(25.7)	(19.4)	(11.9)	(5.6)	(1.5)	(0.9)	(1.3)	(100.0)				
先天性代謝異常による腎不全	2	15	21	25	23	7	8	3	1	2	107	62	169	105.40	37.32
(%)	(1.9)	(14.0)	(19.6)	(23.4)	(21.5)	(6.5)	(7.5)	(2.8)	(0.9)	(1.9)	(100.0)				
腎・尿路結核	1	6	18	15	13	5	7	5	1	71	35	106	109.15	36.52	
(%)	(1.4)	(8.5)	(25.4)	(21.1)	(18.3)	(7.0)	(9.9)	(7.0)	(1.4)	(1.4)	(100.0)				
腎・尿路結石	4	28	50	60	45	35	18	2	1	2	245	139	384	105.33	32.59
(%)	(1.6)	(11.4)	(20.4)	(24.5)	(18.4)	(14.3)	(7.3)	(0.8)	(0.4)	(0.8)	(100.0)				
腎・尿路腫瘍	9	35	91	110	114	33	24	11	4	2	433	211	644	105.71	31.52
(%)	(2.1)	(8.1)	(21.0)	(25.4)	(26.3)	(7.6)	(5.5)	(2.5)	(0.9)	(0.5)	(100.0)				
閉塞性尿路障害	4	25	69	77	68	40	23	4	4	2	316	149	465	107.72	32.22
(%)	(1.3)	(7.9)	(21.8)	(24.4)	(21.5)	(12.7)	(7.3)	(1.3)	(1.3)	(0.6)	(100.0)				
骨髄腫	4	9	20	31	27	20	7	4	3	4	129	84	213	114.59	41.99
(%)	(3.1)	(7.0)	(15.5)	(24.0)	(20.9)	(15.5)	(5.4)	(3.1)	(2.3)	(3.1)	(100.0)				
腎形成不全	2	24	47	50	59	36	20	8	1	2	249	119	368	111.43	34.89
(%)	(0.8)	(9.6)	(18.9)	(20.1)	(23.7)	(14.5)	(8.0)	(3.2)	(0.4)	(0.8)	(100.0)				
不明	202	1,139	2,857	3,236	2,592	1,514	804	354	133	73	12,704	7,774	20,478	107.47	33.02
(%)	(1.6)	(9.0)	(20.9)	(25.5)	(20.4)	(11.9)	(6.3)	(2.8)	(1.0)	(0.6)	(100.0)				
再導入	14	64	162	178	169	103	73	31	6	11	811	353	1,164	111.94	36.43
(%)	(1.7)	(7.9)	(20.0)	(21.9)	(20.8)	(12.7)	(9.0)	(3.8)	(0.7)	(1.4)	(100.0)				
その他	76	314	681	799	688	464	223	118	45	34	3,442	1,964	5,406	109.06	35.16
(%)	(2.2)	(9.1)	(19.8)	(23.2)	(20.0)	(13.5)	(6.5)	(3.4)	(1.3)	(1.0)	(100.0)				
合計	2,545	13,067	28,743	33,990	27,201	16,636	8,533	3,729	1,443	968	136,855	72,681	209,536	107.10	33.62
(%)	(1.9)	(9.5)	(21.0)	(24.8)	(19.9)	(12.2)	(6.2)	(2.7)	(1.1)	(0.7)	(100.0)				
記載なし															
(%)															
総計	2,545	13,067	28,743	33,990	27,201	16,636	8,533	3,729	1,443	968	136,855	72,681	209,536	107.10	33.62
(%)	(1.9)	(9.5)	(21.0)	(24.8)	(19.9)	(12.2)	(6.2)	(2.7)	(1.1)	(0.7)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4623##

(C)Japanese Society for Dialysis Therapy

2016年調査